

Table 2.35-- PREVALENCE OF VARIOUS SUBSTANCES WITH STUDENTS IN THE STATE OF HAWAII, 2007 AND 2011, AND UNITED STATES, 2011

[In percentages]

Substance	Hawaii		U.S.
	2007	2011	2011
Used marijuana in the past 30 days	15.7	21.9	23.1
Ever used any form of cocaine	5.6	6.4	6.8
Glue, aerosol spray can, or paints/spray use 1/	11.4	9.7	11.4
Ever used methamphetamines	4.5	3.4	3.8
Ever used ecstasy	4.6	9.0	8.2
Ever used prescription drugs without a doctor's prescription	(NA)	14.3	20.7
Alcohol use 2/	29.1	29.1	38.7
Binge drinking 3/	14.9	15.4	21.9
Ever tried any form of tobacco	43.7	33.5	(NA)

NA Not available.

1/ Ever sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high.

2/ Had at least one drink of alcohol on at least 1 day in the past 30 days.

3/ Had five or more drinks of alcohol in a row within a couple of hours on at least 1 day in the past 30 days.

Source: Hawaii Department of Education, Hawaii Department of Health, and the University of Hawaii's Curriculum Research & Development Group "*Hawaii Youth Risk Behaviors Hawaii School Health Survey - Survey Results 2011 - State of Hawaii Public High School Report*" and the Hawaii Youth Tobacco Survey.