

Table 20.13-- MOST ABUNDANT AND HEAVIEST FISH SPECIES AT SPECIFIED LOCATIONS: YEAR ENDED JUNE 30, 1998

[Three leading species at each location, based on both individuals per acre and pounds per acre]

Location and species	Individuals per acre		Pounds per acre	
	Rank	Number	Rank	Number
Hanauma Bay MLCD (inshore):				
Manini (<i>Acanthurus triostegus</i>)	1	697	8	68
Saddle-back wrasse (<i>Thalassoma duperrey</i>)	2	450	(1/)	(NA)
Parrotfishes (Family Scaridae)	3	290	(1/)	(NA)
Nenue (<i>Kyphosus</i> sp.)	6	211	1	561
Pualu (<i>Acanthurus xanthopterus</i>)	(1/)	(NA)	2	298
Mullet (<i>Mugil cephalus</i>)	(1/)	(NA)	3	278
Waikiki MLCD:				
Ohua/green wrasse	1	523	6	5
Parrotfishes (Family Scaridae)	2	372	7	5
Saddle-back wrasse (<i>Thalassoma duperrey</i>)	3	247	5	15
Kala (<i>Naso unicornis</i>)	7	77	1	43
Manini (<i>Acanthurus triostegus</i>)	4	209	2	21
Maiiii/lavender tang (<i>Acanthurus nigrofuscus</i>)	5	123	3	18
Kealakekua Bay MLCD:				
Kole (<i>Ctenochaetus strigosus</i>)	1	323	3	30
Yellow tang (<i>Zebrasoma flavescens</i>)	2	312	1	41
Blackfin damsel (<i>Chromis vanderbilti</i>)	3	237	(1/)	(NA)
Ele ele (<i>Melichthys niger</i>)	10	85	2	18

NA Not available.

MLCD Marine Life Conservation District.

1/ Not in first 10.

Source: Hawaii State Department of Land and Natural Resources, Division of Aquatic Resources, *Statewide Marine Research and Surveys, Survey of Fish and Habitat, July 1, 1997 to June 30, 1998.*