

**Table 5.26-- WATER QUALITY AT SELECTED PUBLIC BEACHES:
1998 AND 1999**

Island and beach	Number of samples		Enterococci density 1/	
	1998	1999	1998	1999
Hawaii:				
Hapuna Beach	12	38	1.4	0.5
Kahaluu Beach	12	39	2.7	3.7
Kealakekua Bay (off curio stand)	12	-	1.9	(X)
Hilo Bay (Mooheau Park)	-	-	(X)	(X)
Spencer Beach Park	12	46	2.5	2.6
Maui:				
Kapalua (Fleming) Beach (north)	12	48	0.9	1.0
Kihei (north)	12	-	2.0	(X)
Makena Beach	12	-	0.8	(X)
Seven Pools	9	3	1.5	0.5
Sheraton Kaanapali (shoreline)	12	39	1.2	0.8
Lanai:				
Hulopoe Bay	-	-	(X)	(X)
Molokai:				
Kaunakakai Harbor	-	-	(X)	(X)
Oahu:				
Ala Moana Park (center)	-	4	(X)	9.7
Ewa Beach Park	-	7	(X)	0.8
Haleiwa Beach	23	25	1.6	3.0
Hanauma Bay	46	49	3.4	4.2
Kailua Beach Park	36	50	2.2	1.9
Kuhio Beach	46	49	5.0	8.7
Makaha Beach	-	-	(X)	(X)
Waimea Beach	44	41	2.0	3.2
Kauai:				
Anini Park Pavilion	12	43	1.4	1.6
Kalapaki Beach (middle)	12	50	6.2	4.7
Kekaha (Oomano Point)	12	-	0.9	(X)
Poipu Beach Pavilion	12	43	1.0	1.4
West of Lydgate Park (wading pool)	13	50	1.5	1.2

X Not applicable.

1/ Geometric mean, number per 100 ml. The geometric mean standard for Enterococci density is 7 per 100 ml.

Source: Hawaii State Department of Health, Clean Water Branch, records.