

Table 20.10-- MOST ABUNDANT AND HEAVIEST FISH SPECIES AT SPECIFIED LOCATIONS: YEAR ENDED JUNE 30, 1999

[Three leading species at each location, based on both individuals per acre and pounds per acre]

Location and species	Individuals per acre		Pounds per acre	
	Rank	Number	Rank	Number
Hanauma Bay MLCD (inshore):				
Manini (<i>Acanthurus triostegus</i>)	1	743	3	97
Mamo (<i>Abudefduf abdominalis</i>)	2	530	7	74
Taape (<i>Lutjanus kasmira</i>)	3	244	2	101
Redlipped parrot (<i>Scarus rubroviolaceus</i>)	(1/)	(NA)	1	508
Waikiki MLCD:				
Manini (<i>Acanthurus triostegus</i>)	1	145	4	11
Saddle-back wrasse (<i>Thalassoma duperrey</i>)	2	133	5	7
Mamo (<i>Abudefduf abdominalis</i>)	3	93	(1/)	(NA)
Kala (<i>Naso unicornis</i>)	(1/)	(NA)	1	22
Clown tang (<i>Naso lituratus</i>)	(1/)	(NA)	2	13
Maiii/lavender tang (<i>Acanthurus nigrofuscus</i>)	5	76	3	12
Kealakekua Bay MLCD:				
Agile damsel (<i>Chromis agilis</i>)	1	657	10	5
Kole (<i>Ctenochaetus strigosus</i>)	2	454	2	22
Maiii/lavender tang (<i>Acanthurus nigrofuscus</i>)	3	387	5	10
Yellow tang (<i>Zebrasoma flavescens</i>)	5	270	1	23
Nae nae (<i>Acanthurus olivaceus</i>)	(1/)	(NA)	3	20

NA Not available.

MLCD Marine Life Conservation District.

1/ Not in first 10.

Source: Hawaii State Department of Land and Natural Resources, Division of Aquatic Resources, *Statewide Marine Research and Surveys, Survey of Fish and Habitat, July 1, 1998 to June 30, 1999.*