

**Table 2.35-- PREVALENCE OF VARIOUS SUBSTANCES WITH STUDENTS IN HAWAII, 2013 TO 2017, AND THE UNITED STATES, 2017**

[In percentages. The survey is administered in odd-numbered years to Hawaii public school students in middle school (grade 6-8) and high school (grade 9-12). Does not include charter or special public schools. Based on a sample and subject to sampling variability]

Substance and education stage	Hawaii			U.S.
	2013	2015	2017	2017
Middle school				
Marijuana use 1/	7.5	7.9	7.6	(NA)
Ever used any form of cocaine	2.5	3.2	3.3	(NA)
Ever used inhalants 2/	7.7	6.5	(NA)	(NA)
Ever used heroin	(NA)	(NA)	(NA)	(NA)
Ever used methamphetamines	2.5	2.7	2.1	(NA)
Ever used hallucinogenic drugs 3/	(NA)	(NA)	(NA)	(NA)
Ever used ecstasy	1.9	1.7	1.5	(NA)
Ever took prescription drugs without a doctor's prescription	4.0	4.6	6.8	(NA)
Current drinker 4/	9.5	10.4	11.6	(NA)
Binge drinking 5/	4.4	5.3	5.4	(NA)
Current smoker 6/	5.2	5.0	5.0	(NA)
Electronic vapor product use 7/	(NA)	15.7	15.7	(NA)
High school				
Marijuana use 1/	18.9	19.4	18.1	19.8
Ever used any form of cocaine	6.5	5.7	7.1	4.8
Ever used inhalants 2/	9.2	8.0	(NA)	6.2
Ever used heroin	3.4	3.3	4.5	1.7
Ever used methamphetamines	4.3	3.8	4.8	2.5
Ever used hallucinogenic drugs 3/	6.0	8.1	7.6	6.6
Ever used ecstasy	8.0	8.0	6.0	4.0
Ever took prescription drugs without a doctor's prescription	12.9	12.3	12.2	(NA)
Current drinker 4/	25.2	25.1	24.5	29.8
Binge drinking 5/	12.7	13.4	(NA)	13.5
Current smoker 6/	10.4	9.7	8.1	8.8
Electronic vapor product use 7/	(NA)	25.1	25.5	13.2

Continued on next page.

**Table 2.35-- PREVALENCE OF VARIOUS SUBSTANCES WITH STUDENTS  
IN HAWAII, 2013 TO 2017, AND THE UNITED STATES, 2017 -- Con.**

NA Not available.

1/ Used marijuana one or more times during the 30 days before the survey.

2/ Used inhalants such as glue, aerosol spray can, or some paints or spray to get high.

3/ Hallucinogenic drugs such as LSD, acid, PCP, angel dust, mescaline, or mushrooms.

4/ Had at least one drink of alcohol on at least 1 day during the 30 days before the survey.

5/ For middle school and high school students, had five or more drinks of alcohol in a row within a couple of hours on at least 1 day during the 30 days before the survey. For the U.S. data for high school students in 2017, four or more drinks of alcohol in a row (if they were female) or five or more drinks of alcohol in a row (if they were male), within a couple of hours, on at least 1 day during the 30 days before the survey.

6/ Smoked cigarettes on at least 1 day during the 30 days before the survey.

7/ Electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz) used at least 1 day during the 30 days before the survey.

Source: Hawaii State Department of Education and Hawaii State Department of Health, Hawaii Youth Risk Behavior Survey, State of Hawaii, "Hawaii School Health Survey: Youth Risk Behavior Survey Module for 2013, 2015 and 2017" <<http://hhdw.org/>> accessed June 8, 2018 and Centers for Disease Control and Prevention, "Youth Risk Behavior Surveillance System," <<https://nccd.cdc.gov/Youthonline/App/Default.aspx>> accessed July 3, 2018.