Table 2.35-- PREVALENCE OF VARIOUS SUBSTANCES AMONG STUDENTS IN HAWAII, 2017 TO 2021, AND THE UNITED STATES, 2021

[In percentages. The survey is administered in odd-numbered years to Hawaii public school students in middle school (grade 6-8) and high school (grade 9-12). Does not include charter or special public schools. Based on a sample and subject to sampling variability]

Hawaii			U.S.
2017	2019	2021	2021
11.6	7.0	(NA)	(NA)
5.4	4.6	(NA)	(NA)
5.0	3.9	1.9	(NA)
	17.7	6.7	(NA)
7.6	7.2	(NA)	(NA)
3.3	2.8	0.8	(NA)
2.1	3.7	(NA)	(NA)
1.5	(NA)	(NA)	(NA)
6.8	10.6	6.6	(NA)
24.5	20.4	16.6	22.7
	_		10.5
			3.8
			18.0
18.1	17.2	12.0	15.8
7 1	5.8	2.5	2.5
			1.3
		1.8	1.8
7.6		_	6.5
6.0	4.5	2.7	2.9
12.2	14.4	10.8	12.2
	11.6 5.4 5.0 15.7 7.6 3.3 2.1 1.5 6.8 24.5 12.6 8.1 25.5 18.1 7.1 4.5 4.8 7.6 6.0	2017 2019 11.6 7.0 5.4 4.6 5.0 3.9 15.7 7.2 3.3 2.8 2.1 3.7 1.5 (NA) 6.8 10.6 24.5 10.9 8.1 5.3 25.5 30.6 18.1 17.2 7.1 5.8 4.5 3.6 4.8 4.5 7.6 (NA) 6.0 4.5	2017 2019 2021 11.6 7.0 (NA) 5.4 4.6 (NA) 5.0 3.9 1.9 15.7 17.7 6.7 7.6 7.2 (NA) 3.3 2.8 0.8 2.1 3.7 (NA) 1.5 (NA) (NA) 6.8 10.6 6.6 24.5 20.4 16.6 12.6 10.9 8.3 8.1 5.3 3.0 25.5 30.6 14.8 18.1 17.2 12.0 7.1 5.8 2.5 4.5 3.6 1.8 4.5 3.6 1.8 4.8 4.5 1.8 7.6 (NA) (NA) 6.0 4.5 2.7

NA Not available.

^{1/} Consumed substance on at least 1 day during the 30 days before the survey.

^{2/} Had four or more drinks of alcohol in a row if they were female, or five or more drinks of alcohol in a row if they were male, within a couple of hours, on at least 1 day during the 30 days before the survey.

^{3/} Ever took a prescription pain medicine without a doctor's prescription or different than prescribed use.

^{4/} Hallucinogenic drugs such as LSD, acid, PCP, angel dust, mescaline, or mushrooms.

Source: Centers for Disease Control and Prevention, "Youth Risk Behavior Survey" for Hawaii and the United States https://nccd.cdc.gov/youthonline accessed May 5, 2023