
Instream Flow Standard Assessment Report

Island of Maui

Hydrologic Unit 6008

Kahoma

November 2018

PR-2018-08



State of Hawaii
Department of Land and Natural Resources
Commission on Water Resource Management



COVER

Satellite image of Kahoma Gulch with the Kahoma Stream flowing into the Pacific Ocean, West Maui [Google Earth, 2008].

Note: This report is intended for both print and electronic dissemination and does not include diacritical marks in spelling of Hawaiian words, names, and place names due to problems associated with its use electronically. However, Commission staff has made attempts to include diacritical marks in direct quotations to preserve accuracy.

Table of Contents

1.0	Introduction.....	1
	General Overview	1
	Current Instream Flow Standard.....	1
	Instream Flow Standards	2
	Interim Instream Flow Standard Process.....	3
	Instream Flow Standard Assessment Report.....	4
	Surface Water Hydrologic Units.....	4
	Surface Water Definitions	5
2.0	Unit Characteristics.....	5
	Geology.....	5
	Soils	5
	Rainfall	6
	Solar Radiation.....	8
	Evaporation	8
	Land Use.....	9
	Land Cover	10
	Flood	11
	Drought.....	12
3.0	Hydrology.....	23
	Streams in Hawaii	23
	Ground Water	25
	Wells in the Kahoma Hydrologic Unit.....	26
	Groundwater Pumping and Salinity Levels	26
	Kahoma and Kanaha Streamflow Characteristics.....	29
	Long-term trends in flow.....	32
4.0	Maintenance of Fish and Wildlife Habitat.....	36
	Hawaii Stream Assessment.....	37
5.0	Outdoor Recreational Opportunities	40
6.0	Maintenance of Ecosystems.....	42
7.0	Aesthetic Values.....	50
8.0	Navigation	51
9.0	Instream Hydropower Generation.....	51
10.0	Maintenance of Water Quality	52
11.0	Conveyance of Irrigation and Domestic Water Supplies	57
12.0	Protection of Traditional and Customary Hawaiian Rights.....	58
	Taro Production	66
	Fishponds.....	71
13.0	Public Trust Uses of Water.....	74
	Hawaiian Home Lands.....	74
	Domestic Water Supply.....	75

14.0	Noninstream Uses	77
	Water Leaving the Kahoma Hydrologic Unit in Ditch Systems	77
	Modifications of Ditch Systems and Groundwater Recharge	84
	Utilization of Important Agricultural Lands.....	85
	Irrigation Needs of the Kahoma Hydrologic Unit	86
15.0	Bibliography.....	90
16.0	Appendices	99

List of Figures

Figure 1-1. Information to consider in setting measurable instream flow standards.....	2
Figure 1-2. Simplified representation of the interim instream flow standard and permanent instream flow standard processes. Keys steps of the adopted interim IFS process are depicted in the left column by the boxes drawn with dotted lines.....	3
Figure 1-3. Quickbird satellite imagery of the Kahoma hydrologic unit and streams in West Maui, Hawaii. (Source: State of Hawaii, Planning Department, 2004; State of Hawaii, Commission on Water Resource Management, 2015c; State of Hawaii, Division of Aquatic Resources, 2005)	7
Figure 1-4. Elevation range of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2004e; U.S. Geological Survey, 2001).....	7
Figure 1-5. USGS topographic map of Kahoma hydrologic unit, Maui, Hawaii. (Source: U.S. Geological Survey, 1996).....	3
Figure 1-6. Major and minor roads in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning 2018).....	4
Figure 2-1. Orographic precipitation in the presence of mountains higher than 6,000 feet.	7
Figure 2-2. Generalized geology of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Sherrod et al., 2007).....	14
Figure 2-3. Soil classification of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015m).....	15
Figure 2-4. Mean annual rainfall of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Giambelluca et al., 2013).....	15
Figure 2-5. Mean annual solar radiation of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015c).....	17
Figure 2-6. Mean annual potential evapotranspiration (Penman-Monteith method) of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Giambelluca et al., 2014)	18
Figure 2-7. State land use district boundaries of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015d).....	19
Figure 2-8. C-CAP land cover of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015k).....	20
Figure 2-9. Hawaii GAP land cover classes of the Kahoma hydrologic unit, Maui, Hawaii (Source: USGS, 2011).	21
Figure 2-10. FEMA flood zone regions in Kahoma hydrologic unit (Source: Federal Emergency Management Agency, 2014).....	22
Figure 3-1. Conceptual diagram illustrating surface water-groundwater interactions in West Maui. (Source: Oki et al., 2010),.....	24
Figure 3-2. Well locations and numbers in the Kahoma hydrologic unit, which spans the Launiupoko and Honokowai aquifer sectors, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b).....	27
Figure 3-3. Monthly reported pumpage, 12-month moving average (12-MAV) and sustainable yield for the Launiupoko Aquifer sector. (Source: Source: State of Hawaii, Commission on Water Resource Management, 2018b).....	28
Figure 3-4. Elevation to freshwater lense, brackish water transition and sea water in the Mahinahina Deep Monitoring Well, West Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018a).....	29
Figure 3-5. Monthly groundwater pumpage and chloride concentration from two municipal (A & B) and one private (C) well in West Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b).	30

Figure 3-6. USGS surface water gage locations by number for the Kahoma hydrologic Unit, Maui, Hawaii. (Source: USGS, 2011).	33
Figure 3-7. Seepage run results and median flow in cubic feet per second from partial record gaging stations of the Kahoma hydrologic unit (Source: State of Hawaii, Commission on Water Resource Management, 2015a; Cheng, 2014).	34
Figure 3-8. Cumulative departures of monthly mean flow from the mean of the monthly flows, Hawaii. This data is based on complete water years from 1913 through 2002. (Oki, 2004, Figure 4)	35
Figure 4-1. Elevational profile of a terminal-estuary stream on the Big Island of Hawaii (Hakalau Stream). (Source: McRae, 2007, adapted from Nishimoto and Kuamoo, 1991 [with permission])	36
Figure 4-2. Classification of stream reaches based on habitat suitability for native species. (Source: Tingley III et al. 2018).	39
Figure 5-1. Public hunting areas for game mammals and coastal resources in Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 1999, 2002b, 2002c, 2004a).	41
Figure 6-1. Simplified ecosystem illustrated in a Hawaiian stream. (Source: Ziegler, 2002, illustration by Keith Kruger).	42
Figure 6-2. Reserves and wetlands for the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2007b; 2015n)	47
Figure 6-3. Distribution of native and alien plant species, and critical habitat for plant and bird species in the Kahoma hydrologic unit. (Source: Scott et al., 1986; State of Hawaii, Office of Planning, 1996; 2004b).	47
Figure 6-4. Density of threatened and endangered plants in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015h)	49
Figure 10-1. Water quality standards for the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015e; 2008). The classifications are general in nature and should be used in conjunction with Hawaii Administrative Rules, Chapter 11-54, Water Quality Standards.	56
Figure 12-1. Generalized process for determining appurtenant water rights. This process is generalized and may not fully explain all possible situations. It does not apply to Hawaiian Homes Lands. If you are Native Hawaiian you may have other water rights.	61
Figure 12-2. Land Commission Awards and associated tax map key parcel numbers for the Kahoma hydrologic unit, Maui.	62
Figure 12-3. Traditional ahupuaa boundaries in the Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Office of Planning, 2015j).	70
Figure 12-4. Tax map key boundaries in the Kahoma hydrologic unit with specific parcels along Kanaha Stream (insert), Maui.	73
Figure 13-1. Hawaiian Home Lands development parcels and other major land owners, West Maui. (Source: State of Hawaii, Department of Hawaiian Home Lands, 2011).	75
Figure 13-2. Water supply service systems for the Lahaina Aquifer System, Maui. (Source: Maui County, 2018).	76
Figure 14-1. Registered monthly average mean daily flow (million gallons per day, mgd) diverted from Kanaha Stream in the Kahoma Hydrologic Unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018c).	78
Figure 14-2. All registered diversions (ID) and ditches identified in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2004d; State of Hawaii, Commission on Water Resource Management, 2015g).	83
Figure 14-3. Mean daily flow in Kahoma Ditch from Kahoma Stream and Lahainaluna Ditch from Kanaha Stream in the Kahoma hydrologic unit. (Source: State of Hawaii, Commission on Water Resource Management, 2018c).	84

Figure 14-4. Estimated recharge for six historical periods between 1926 and 2004, central and west Maui, Hawaii. (Source: Engott and Vana, 2007).....85

Figure 14-5. Agricultural Lands of Importance designated in the Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Office of Planning, 2015g).....88

Figure 14-6. 2015 Baseline Agricultural Land Use Map for the Kahoma hydrologic Unit, Maui. (Source: (Perroy et al. 2016).....89

List of Tables

Table 2-1. Area and percentage of surface geologic features for Kahoma hydrologic unit, Maui, Hawaii. (Source: Sherrod et al, 2007).....	5
Table 2-2. Area and percentage of soil types for the Kahoma hydrologic unit, Maui, Hawaii. (Source: USDA, 1986).....	6
Table 2-3. Fog drip to rainfall ratios for the windward slopes of Mauna Loa on the island of Hawaii and approximate contributions to the Kahoma Hydrologic Unit based on an elevation range of 2000-5780 ft.....	8
Table 2-4. C-CAP land cover classes and area distribution in Kahoma hydrologic unit, Maui, Hawaii. (Source: National Oceanographic and Atmospheric Agency, 2015).....	10
Table 2-5. HI-GAP land cover classes and area distribution in Kahoma hydrologic unit, Maui, Hawaii. (Source: USGS, 2011).....	11
Table 2-6. Drought risk areas for Maui. (Source: University of Hawaii, 2003).....	12
Table 3-1. Selected discharge exceedance values from historic stream gaging data in the Kahoma hydrologic unit, Maui, Hawaii. (Source: USGS NWIS, 2018). [in cubic feet per second (million gallons per day)].....	25
Table 3-2. Information of wells located in Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b). [IRRLA = Landscape irrigation; MUNCO = Municipal, County; DOM = domestic; UNU = unused; AGRCP = Agriculture row crop; HHFDC = Hawaii Housing Finance & Development Corporation).....	28
Table 3-3. Selected natural-flow duration discharge exceedance values for the current (1984-2013) climate period based on partial record gaging station for the Kahoma hydrologic unit, Maui, Hawaii. (Source: Cheng 2014) [Flows are in cubic feet per second (million gallons per day)].....	31
Table 3-4. Discharge measurements at gage 6-123 and estimated diversion amounts from diversion 951 on Kahoma Stream. (Source: CWRM) [Flows are in cubic feet per second (million gallons per day)].....	31
Table 4-1. List of commonly mentioned native stream organisms. (Source: State of Hawaii, Division of Aquatic Resources, 1993).....	36
Table 4-2. Size distribution of <i>Awaous stamineus</i> by habitat type in Kahoma Stream.....	38
Table 6-1. Hawaii Stream Assessment indicators of riparian resources for Kahoma hydrologic unit, Maui, Hawaii.	43
Table 6-2. Management areas located within Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Division of Forestry and Wildlife, 2008a).....	44
Table 6-3. Watershed partnerships associated with the Kahoma hydrologic unit. (Source: State of Hawaii, Division of Forestry and Wildlife, 2008b).....	44
Table 6-4. Wetland classifications for Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015n).....	45
Table 6-5. Density of threatened and endangered plants for Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015i).....	45
Table 6-6. Estimated Net Present Value (NPV) for Koolau [Oahu] Forest Amenities. (Source: Kaiser, B. et al., n.d.).....	46
Table 12-1. Summary of water use calculated from loi and loi complexes by island, and the entire state. (Source: Gingerich et al., 2007, Table 10).....	67
Table 12-2. Water-temperature statistics based on measurements collected at 15-minute intervals for loi complexes on the island of Maui. (Source: Gingerich et al., 2007, Table 7).....	68
Table 12-3. Cultural resource elements evaluated as part of the Hawaii Stream Assessment for Kahoma hydrologic unit, Maui.....	72

Table 13-1. Current total and domestic (single family) water use by subdistrict for the Lahaina-Napili service area provided by the Maui County Department of Water Supply. (Source: Maui County, 2018)	76
Table 14-1. Surface water registrations for the Kahoma Hydrologic Unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2015e).....	78
Table 14-2. Registered diversions in the Kahoma hydrologic unit.....	79
Table 14-3. Selected off-stream water use statistics in the Kahoma hydrologic unit, Maui, Hawaii. (Source: CWRM, 2018b).....	84
Table 14-4. Agricultural Lands of Importance to the State of Hawaii and area distributions in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015g).....	86
Table 14-5. Mean drip irrigation water demand estimates for three crops in the Kahoma service area based on IWREDSS scenarios modeled for two row crop management and two soil management techniques given a 50 ft depth to water table. Demand is based on various drought scenarios.	87

Acronyms and Abbreviations

A&B	Alexander & Baldwin
AG	agricultural
ALISH	Agricultural Lands of Importance to the State of Hawaii
ALUM	agricultural land use maps [prepared by HDOA]
BFQ	base flow statistics
BLNR	Board of Land and Natural Resources (State of Hawaii)
C-CAP	Coastal Change Analysis Program
cfs	cubic feet per second
Code	State Water Code (State of Hawaii)
COM	commercial
Commission	Commission on Water Resource Management (DLNR)
CPRC	Compilation of Public Review Comments (PR-2008-07, CWRM)
CWA	Clean Water Act (EPA)
CWRM	Commission on Water Resource Management (State of Hawaii)
DAR	Division of Aquatic Resources (State of Hawaii)
DHHL	Department of Hawaiian Home Lands (State of Hawaii)
DLNR	Department of Land and Natural Resources (State of Hawaii)
DOH	Department of Health (State of Hawaii)
DWS	Department of Water Supply (County of Maui)
EA	Environmental Assessment
EIS	Environmental Impact Statement
EPA	United States Environmental Protection Agency
FEMA	Federal Emergency Management Agency (Department of Homeland Security)
FILEREF	File Reference [in the Commission's records of registered diversions]
ft	feet
gad	gallons per acre per day
GIS	Geographic Information Systems
G.L.	Government Lease
GOV	government
gpm	gallons per minute
Gr.	Grant
HAR	Hawaii Administrative Rules
HC&S	Hawaiian Commercial and Sugar Company
HDOA	State Department of Agriculture (State of Hawaii)
HI-GAP	Hawaii Gap Analysis Program
HOT	hotel
HSA	Hawaii Stream Assessment
IFS	instream flow standard
IFSAR	Instream Flow Standard Assessment Report
IND	industry
IRR	irrigation requirements
IWREDSS	Irrigation Water Requirement Estimation Decision Support System
KAA	Kekaha Agriculture Association
KIUC	Kauai Island Utility Cooperative
LCA	Land Commission Award
LUC	Land Use Commission (State of Hawaii)
MECO	Maui Electric Company
MF	multi-family residential
mgd	million gallons per day
Mgal/d	million gallons per day
mi	mile
MOU	Memorandum of Understanding

na	not available
NAWQA	National Water Quality Assessment (USGS)
NHLC	Native Hawaiian Legal Corporation
NIR	net irrigation requirements
NPDES	National Pollutant Discharge Elimination System
NPV	Net Present Value
NRCS	Natural Resource Conservation Service (USDA)
NVCS	National Vegetation Classification System
por.	Portion
REL	religious
RMT	R.M. Towill Corporation
SCS	Soil Conservation Service (United States Department of Agriculture) Note: The SCS is now called the Natural Resources Conservation Service (NRCS)
SF	single family residential
SPI	Standardized Precipitation Index
sq mi	square miles
TFQ	total flow statistics
TFQ ₅₀	50 percent exceedence probability
TFQ ₉₀	90 percent exceedence probability
TMDL	Total Maximum Daily Load
TMK	Tax Map Key
UHERO	University of Hawaii's Economic Research Organization
USDA	United States Department of Agriculture
USFWS	United States Fish and Wildlife Service (Department of the Interior)
USGS	United States Geological Survey (Department of the Interior)
WQS	Water Quality Standards
WRPP	Water Resource Protection Plan (Commission on Water Resource Management)
WTF	water treatment facility

1.0 Introduction

General Overview

The hydrologic unit of Kahoma is located in West Maui, on the the western flank of Puu Kukui Mountain which forms the western part of the Hawaiian island of Maui (Figure 1-3). It covers an area of 5.79 square miles from 5,764 feet elevation to the sea with a mean basin elevation of 2,390 feet and a mean basin slope of 71.2 percent (Figure 1-4 and 1-5). Sixty-nine percent of the basin has a slope greater than 30 percent. The longest flow path in Kahoma is 7.8 miles in length, traversing in a westernly direction from its headwaters to Lahaina Harbor. The Kahoma Stream is composed of two main branches, Kahoma and Kanaha streams. Kahoma Stream has one main tributary, Halona Stream, which only flows intermittantly. A number of smaller tributaries flow only intermittently in response to rainfall-runoff. Baseflow is supported by dike-impounded groundwater in the upper reaches with two development tunnels built to augment spring flow. The USGS estimated that the Kahoma branch would flow continuously from the current diversion to the confluence with Kanaha 85 to 90 percent of the time while the Kanaha branch is estimated to flow at least 95 percent of the time to the confluence. However, following voluntary restoration of streamflow by the landowner Kamehameha Schools Bishop Estate (KSBE) in 2010, continuous streamflow has reached the confluence for many years. Downstream of the confluence, Kahoma Stream is estimated to flow to the ocean naturally 95 percent of the time, but this is influenced by the concrete lined channel which limits seepage loss. The basin has a mean annual precipitation of 102 inches. The highest elevation sections of the hydrologic unit are made up of conservation land owned by KSBE and the State of Hawaii Department of Land and Natural Resources (DLNR). The City of Lahaina, a census designated place that includes Kaanapali and Kapalua beach resorts, is located along the coast with a total population of 11, 704 people (U.S. Census Bureau Office of Planning 2011). The lower elevations are very dry and occupied by grasses, shrubs and alien trees, while the upper regions support native ohia- and koa-dominated forests. In the mid elevation reaches along Kahoma Stream there are cultivated taro loi re-established following stream restoration, while along Kanaha Stream there is currently insufficient water to support the historic loi due to the stream diversion. Two primary state roads exist in the region, but a number of secondary roads provide access to neighborhoods, as well as commercial and industrial areas of the hydrologic unit. Former agricultural roads provide limited access to large areas of the hydrologic unit (Figure 1-6). Two different irrigation ditch systems diverted water: one from the Kahoma branch and one from the Kanaha branch, which continues to be used for municipal water supply and agriculture irrigation.

Current Instream Flow Standard

The current interim instream flow standard (IFS) for Kahoma Stream was established by way of Hawaii Administrative Rules (HAR) §13-169-44, which, in pertinent part, reads as follows:

Interim instream flow standard for Maui. The Interim Instream Flow Standard for all streams on Maui, as adopted by the commission on water resource management on June 15, 1988, shall be that amount of water flowing in each stream on the effective date of this standard, and as that flow may naturally vary throughout the year and from year to year without further amounts of water being diverted offstream through new or expanded diversions, and under the stream conditions existing on the effective date of the standard.

The current interim IFS became effective on December 31, 1988. Streamflow was not measured on that date; therefore, the current interim IFS is not a quantifiable value.


Instream Flow Standards

Under the State Water Code (Code), Chapter 174C, Hawaii Revised Statutes (HRS), the Commission on Water Resource Management (Commission) has the responsibility of establishing IFS on a stream-by-stream basis whenever necessary to protect the public interest in the waters of the State. Early in its history, the Commission recognized the complexity of establishing IFS for the State’s estimated 376 perennial streams and instead set interim IFS at “status quo” levels. These interim IFS were defined as the amount of water flowing in each stream (with consideration for the natural variability in stream flow and conditions) at the time the administrative rules governing them were adopted in 1988 and 1989.

The Hawaii Supreme Court, upon reviewing the Waiahole Ditch Contested Case Decision and Order, held that such “status quo” interim IFS were not adequate to protect streams and required the Commission to take immediate steps to assess stream flow characteristics and develop quantitative interim IFS for affected Windward Oahu streams, as well as other streams statewide. The Hawaii Supreme Court also emphasized that “instream flow standards serve as the primary mechanism by which the Commission is to discharge its duty to protect and promote the entire range of public trust purposes dependent upon instream flows.”

To the casual observer, IFS may appear relatively simple to establish upon a basic review of the Code provisions. However, the complex nature of IFS becomes apparent upon further review of the individual components that comprise surface water hydrology, instream uses, noninstream uses, and their interrelationships. The Commission has the distinct responsibility of weighing competing uses for a limited resource in a legal realm that is continuing to evolve. The following illustration (Figure 1-1) was developed to illustrate the wide range of information, in relation to hydrology, instream uses, and noninstream uses that should be addressed in conducting a comprehensive IFS assessment.

Figure 1-1. Information to consider in setting measurable instream flow standards.




Interim Instream Flow Standard Process

The Code provides for a process to amend an interim IFS in order to protect the public interest pending the establishment of a permanent IFS. The Code, at §174C-71(2), describes this process including the role of the Commission to “weigh the importance of the present or potential instream values with the importance of the present or potential uses of water for noninstream purposes, including the economic impact of restricting such uses.”

Recognizing the complexity of establishing measurable IFS, while cognizant of the Hawaii Supreme Court’s mandate to designate interim IFS based on best available information under the Waiahole Combined Contested Case, the Commission at its December 13, 2006 meeting authorized staff to initiate and conduct public fact gathering. Under this adopted process (reflected in the left column of Figure 1-2), the Commission staff will conduct a preliminary inventory of best available information upon receipt of a petition to amend an existing interim IFS. The Commission staff shall then seek agency review and comments on the compiled information (compiled in an Instream Flow Standard Assessment Report) in conjunction with issuing a public notice for a public fact gathering meeting. Shortly thereafter (generally within 30 days), the Commission staff will conduct a public fact gathering meeting in, or near, the hydrologic unit of interest.

Figure 1-2. Simplified representation of the interim instream flow standard and permanent instream flow standard processes. Keys steps of the adopted interim IFS process are depicted in the left column by the boxes drawn with dotted lines.



Instream Flow Standard Assessment Report

The Instream Flow Standard Assessment Report (IFSAR) is a compilation of the hydrology, instream uses, and noninstream uses related to a specific stream and its respective surface water hydrologic unit. The report is organized in much the same way as the elements of IFS are depicted in Figure 1-1. The purpose of the IFSAR is to present the best available information for a given hydrologic unit. This information is used to determine the interim IFS recommendations, which is compiled as a separate report. The IFSAR is intended to act as a living document that should be updated and revised as necessary, thus also serving as a stand-alone document in the event that the Commission receives a subsequent petition solely for the respective hydrologic unit.

Each report begins with an introduction of the subject hydrologic unit and the current IFS status. Section 2.0 is comprised of the various hydrologic unit characteristics that, both directly and indirectly, impact surface water resources. Section 3.0 contains a summary of available hydrologic information, while Sections 4.0 through 12.0 summarize the best available information for the nine instream uses as defined by the Code. Section 13.0 describes public trust uses of water not covered in other sections. Noninstream uses are summarized in Section 14.0. Maps are provided at the end of each section to help illustrate information presented within the section's text or tables. Finally, Section 15.0 provides a comprehensive listing of cited references and is intended to offer readers the opportunity to review IFSAR references in further detail.

An important component of the IFSAR and the interim IFS process is the Compilation of Public Review Comments (CPRC). The CPRC serves as a supporting document containing the oral and written comments that are submitted as part of the initial public review process. Comments referred to within the IFSAR will identify both the section and page number where the original comment can be located in the CPRC. For example, a reference to "8.0-3" indicates the third page of comments in Section 8.0 of the CPRC.

Following the preparation of the IFSAR and initial agency and public review, information may be added to the IFSAR at any time. Dates of revision will be reflected as such. Future review of the IFSAR, by agencies and the public, will only be sought when a new petition to amend the interim (or permanent) instream flow standard is pending. Recommendations for IFS amendments are prepared separately as a stand-alone document. Thus, the IFSAR acts solely as a compendium of best available information and may be revised further without the need for subsequent public review following its initial preparation.

Surface Water Hydrologic Units

Early efforts to update the Commission's Water Resource Protection Plan (WRPP) highlighted the need for surface water hydrologic units to delineate and codify Hawaii's surface water resources. Surface water hydrologic units served as an important first-step towards improving the organization and management of surface water information that the Commission collects and maintains, including diversions, stream channel alterations, and water use.

In developing the surface water hydrologic units, the Commission staff reviewed various reports to arrive at a coding system that could meet the requirements for organizing and managing surface water information in a database environment, and could be easily understood by the general public and other agencies. For all intents and purposes, surface water hydrologic units are synonymous with watershed areas. Though Commission staff recognized that while instream uses may generally fall within a true surface drainage area, noninstream uses tend to be land-based and therefore may not always fall within the same drainage area.

In June 2005, the Commission adopted the report on surface water hydrologic units and authorized staff to implement its use in the development of information databases in support of establishing IFS (State of

Hawaii, Commission on Water Resource Management, 2005a). The result is a surface water hydrologic unit code that is a unique combination of four digits. This code appears on the cover of each IFSAR above the hydrologic unit name.

Surface Water Definitions

Listed below are the most commonly referenced surface water terms as defined by the Code.

Agricultural use. The use of water for the growing, processing, and treating of crops, livestock, aquatic plants and animals, and ornamental flowers and similar foliage.

Channel alteration. (1) To obstruct, diminish, destroy, modify, or relocate a stream channel; (2) To change the direction of flow of water in a stream channel; (3) To place any material or structures in a stream channel; and (4) To remove any material or structures from a stream channel.

Continuous flowing water. A sufficient flow of water that could provide for migration and movement of fish, and includes those reaches of streams which, in their natural state, normally go dry seasonally at the location of the proposed alteration.

Domestic use. Any use of water for individual personal needs and for household purposes such as drinking, bathing, heating, cooking, noncommercial gardening, and sanitation.

Ground water. Any water found beneath the surface of the earth, whether in perched supply, dike-confined, flowing, or percolating in underground channels or streams, under artesian pressure or not, or otherwise.

Hydrologic unit. A surface drainage area or a ground water basin or a combination of the two.

Impoundment. Any lake, reservoir, pond, or other containment of surface water occupying a bed or depression in the earth's surface and having a discernible shoreline.

Instream Flow Standard. A quantity of flow of water or depth of water which is required to be present at a specific location in a stream system at certain specified times of the year to protect fishery, wildlife, recreational, aesthetic, scenic, and other beneficial instream uses.

Instream use. Beneficial uses of stream water for significant purposes which are located in the stream and which are achieved by leaving the water in the stream. Instream uses include, but are not limited to:

- (1) Maintenance of fish and wildlife habitats;
- (2) Outdoor recreational activities;
- (3) Maintenance of ecosystems such as estuaries, wetlands, and stream vegetation;
- (4) Aesthetic values such as waterfalls and scenic waterways;
- (5) Navigation;
- (6) Instream hydropower generation;
- (7) Maintenance of water quality;
- (8) The conveyance of irrigation and domestic water supplies to downstream points of diversion; and
- (9) The protection of traditional and customary Hawaiian rights.

Interim instream flow standard. A temporary instream flow standard of immediate applicability, adopted by the Commission without the necessity of a public hearing, and terminating upon the establishment of an instream flow standard.

Municipal use. The domestic, industrial, and commercial use of water through public services available to persons of a county for the promotion and protection of their health, comfort, and safety, for the protection of property from fire, and for the purposes listed under the term "domestic use."

Noninstream use. The use of stream water that is diverted or removed from its stream channel and includes the use of stream water outside of the channel for domestic, agricultural, and industrial purposes.

Reasonable-beneficial use. The use of water in such a quantity as is necessary for economic and efficient utilization, for a purpose, and in a manner which is both reasonable and consistent with the state and county land use plans and the public interest.

Stream. Any river, creek, slough, or natural watercourse in which water usually flows in a defined bed or channel. It is not essential that the flowing be uniform or uninterrupted. The fact that some parts of the bed or channel have been dredged or improved does not prevent the watercourse from being a stream.

Stream channel. A natural or artificial watercourse with a definite bed and banks which periodically or continuously contains flowing water. The channel referred to is that which exists at the present time, regardless of where the channel may have been located at any time in the past.

Stream diversion. The act of removing water from a stream into a channel, pipeline, or other conduit.

Stream reach. A segment of a stream channel having a defined upstream and downstream point.

Stream system. The aggregate of water features comprising or associated with a stream, including the stream itself and its tributaries, headwaters, ponds, wetlands, and estuary.

Surface water. Both contained surface water--that is, water upon the surface of the earth in bounds created naturally or artificially including, but not limited to, streams, other watercourses, lakes, reservoirs, and coastal waters subject to state jurisdiction--and diffused surface water--that is, water occurring upon the surface of the ground other than in contained water bodies. Water from natural springs is surface water when it exits from the spring onto the earth's surface.

Sustainable yield. The maximum rate at which water may be withdrawn from a water source without impairing the utility or quality of the water source as determined by the Commission.

Time of withdrawal or diversion. In view of the nature, manner, and purposes of a reasonable and beneficial use of water, the most accurate method of describing the time when the water is withdrawn or diverted, including description in terms of hours, days, weeks, months, or physical, operational, or other conditions.

Watercourse. A stream and any canal, ditch, or other artificial watercourse in which water usually flows in a defined bed or channel. It is not essential that the flowing be uniform or uninterrupted.

Figure 1-3. Quickbird satellite imagery of the Kahoma hydrologic unit and streams in West Maui, Hawaii. (Source: State of Hawaii, Planning Department, 2004; State of Hawaii, Commission on Water Resource Management, 2015c; State of Hawaii, Division of Aquatic Resources, 2005)





Figure 1-4. Elevation range of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2004e; U.S. Geological Survey, 2001)



Prepared by the Department of Land and Natural Resources,
 Commission on Water Resource Management
 Transverse Mercator Projection, Zone 4, North American Datum 1983

Figure 1-5. USGS topographic map of Kahoma hydrologic unit, Maui, Hawaii. (Source: U.S. Geological Survey, 1996)





Figure 1-6. Major and minor roads in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning 2018)



2.0 Unit Characteristics

Geology

Kahoma is on the western flanks of Puu Kukui, the West Maui volcano (Figure 2-2). There were three separate volcanic phases in West Maui (Stearns and Macdonald 1942). The first phase, called the Wailuku Volcanic Series, makes up about 97% of the volume of the volcano, and consists of thin pahoehoe and aa flows of tholeiitic olivine basalt with minor plagioclase basalt as thick as 1700 meters (Table 2-1). The summit of this shield building phase eventually collapsed to form a caldera about 3.2 km across, with horizontal post-caldera lava flows. The Honolua Volcanic Series, the second phase, is dominated by alkali rocks that formed an incomplete cap to the volcano ranging from single flows less than 10 m thick to several flows as thick as 230 m on the northeastern slopes (Macdonald and Abbott 1970). Occasionally viscous trachytic magma formed domes with steeply sloping flow planes, especially in the Kahoma and Launiupoko areas. Many dike and vent formations were produced in the Honolua Series although due to the lack of well-developed rift zones, their distribution is somewhat irregular. The Honolua Series finished about 500 kya, followed by a brief third phase of activity which included four small eruptions occurring on the southwestern slope forming the Lahaina Volcanic Series. These lava flows were all silica undersaturated basanitoids or basanites. The rapid erosion and valley incision of West Maui has produced broad alluvial fans with unconsolidated dunes of lithified to semi-lithified calcareous sand on the western slopes. The generalized geology of the Kahoma hydrologic unit is depicted in Figure 2-2.

Table 2-1. Area and percentage of surface geologic features for Kahoma hydrologic unit, Maui, Hawaii. (Source: Sherrod et al, 2007)

Name	Rock Type	Area (mi ²)	Percent (%)
Wailuku Volcanics	Intrusive rocks, Caldera complex, Dike complex, Pahoehoe and Aa lava flows	4.519	78.0
Lahaina Volcanics	Lava flows	0.755	13.0
Alluvium	Sand and Gravel	0.307	5.3
Older alluvium	Lithified sand and gravel	0.108	1.9
Honolua Volcanics	Pahoehoe and Aa, Conglomerate	0.102	1.8

Soils

The U.S. Department of Agriculture's Natural Resources Conservation Service (formerly known as the Soil Conservation Service) divides soils into hydrologic soil groups (A, B, C, and D) according to the rate at which infiltration (intake of water) occurs when the soil is wet. The higher the infiltration rate, the faster the water is absorbed into the ground and the less there is to flow as surface runoff. Group A soils have the highest infiltration rates; group D soils have the lowest. In the Kahoma hydrologic unit, soils are largely dominated by Wahikuli Series and Lahaina Series, with 16.3% of soils in hydrologic soil Group B, 30.9% of soils in Group C and 52.0% of soils in Group D (Table 2-2). Most of the unit consists of soils in Group B, silty loam or loam with moderate infiltration rates, or Group D, clay loam, silty clay loam, or clay, with high runoff potential. The less sloping tops of ridges and interfluvies (regions of higher land between valleys in the same hydrologic unit) are poorly drained but still support moderate infiltration. The lowlands are composed of well-drained clay and rough broken land – very steep land broken by numerous intermittent drainage channels (Figure 2-3). The numerous gulches and mountainsides have slopes of 40 to 70 percent, leading to high rates of slopewash. Runoff is rapid in these soils and geologic erosion is active. The soils of rough broken land are not uniform. The clay is moderately permeable with slow to medium runoff and a slight to moderate erosion hazard (U.S. Department of Agriculture, National Resource Conservation Service, 1986).

Table 2-2. Area and percentage of soil types for the Kahoma hydrologic unit, Maui, Hawaii. (Source: USDA, 1986)

Soil Series Unit	Area (mi ²)	Percent (%)
Beaches	0.003	0.05%
Wainee	0.018	0.31%
Ewa	0.187	3.24%
Lahaina	0.649	11.21%
Pulehu	0.106	1.82%
Alaeloa	0.088	1.51%
Rough broken land	0.296	5.11%
Wahikuli	1.403	24.23%
Olelo	0.255	4.40%
Rock land	0.166	2.87%
Rock outcrop	0.514	8.87%
Rough broken & stony land	0.022	0.38%
Rough mountainous land	2.057	35.53%
Water	0.028	0.49%

The Wainee soils are deep, extremely stony silty clay, well drained, and weathered from basic igneous rock on alluvial fans of 3-15 percent slope (U. S. Department of Agriculture, Natural Resources Conservation Service, Conservation Engineering Division, 1986). These soils are excellent for agriculture. The center of the hydrologic unit consists of rough broken and rough mountainous soils while alluvium (riverwash) dominates the lower reaches (Figure 2-3).

Rainfall

The West Maui Mountains are the driving force affecting the distribution of rainfall in Kahoma with rainfall affected by both the orographic¹ effect and the rain shadow effect (Figure 2-1). Orographic precipitation occurs when the prevailing northeasterly trade winds lift warm air up the windward side of the mountains into higher elevations where cooler temperatures persist. As moist air cools, water condenses and the air mass releases precipitation. As a result, frequent and heavy rainfall is observed at the windward mountain slopes. The temperature inversion zone typically extends from 6,560 feet to 7,874 feet. This region is influenced by a layer of moist air below and dry air above, making climate extremely variable (Giambelluca and Nullet, 1992). The fog drip zone occurs below the elevation where cloud height is restricted by the temperature inversion, where temperature increases with elevation (Sholl et al., 2002). Fog drip is a result of cloud-water droplets impacting vegetation (Scholl et al, 2002) and can contribute significantly to ground water recharge. Above the inversion zone, the air is dry and the sky is frequently clear (absence of clouds) with high solar radiation, creating an arid atmosphere with little rainfall.


A majority of the mountains in Hawaii peak in the fog drip zone, where cloud-water is intercepted by vegetation. In such cases, air passes over the mountains, warming and drying while descending on the leeward mountain slopes. Precipitation on Puu Kukui is influenced by its position relative to the trade winds. The highest position lies in the cloudy layer below the trade wind inversion resulting in peak rainfall for the region. Mean annual rainfall on Puu Kukui measured by the USGS (station 380.0) and Maui Land & Pineapple since 1928 is about 362 inches. The fog drip zone on the windward side of

¹ Orographic refers to influences of mountains and mountain ranges on airflow, but also used to describe effects on other meteorological quantities such as temperature, humidity, or precipitation distribution.

islands extends from the cloud base level at 1,970 feet to the lower limit of the most frequent temperature inversion base height at 6,560 feet (Giambelluca and Nullet, 1992). The steep gradient around the island forces moisture-laden air to rapidly rise in elevation (over 3,000 feet) in a short distance, resulting in a rapid release of rainfall in one location. Finally, the relatively round, conical shape of the West Maui Mountains exposes all sides of the peak to wind and moisture.

The Kahoma hydrologic unit is situated on the leeward flank of the West Maui Mountains and as such receives less orographic rainfall than windward slopes and is impacted by the rain shadow effect. Wind will blow rainfall over the interfluve from east/northeast facing hydrologic units, contributing to higher rainfall in the upper elevations (Figure 2-4). The high spatial variability in rainfall is evident by the large variation in mean annual rainfall across the hydrologic unit. For the whole hydrologic unit, mean annual rainfall averages just 81.4 inches. Above 2000 ft, rainfall is highest during the months of December and January, where the mean monthly rainfall is 13.1 inches and 13.8 inches, respectively. Irregular southerly Kona storm systems may produce localized intense rainfall on southeast facing slopes.

Figure 2-1. Orographic precipitation in the presence of mountains higher than 6,000 feet.



Currently, fog drip data for leeward slopes are non-existent. Shade (1999) used the monthly fog drip to rainfall ratios for the windward slopes of Mauna Loa on the island of Hawaii (Table 2-3) to calculate fog drip contribution to the water-budget in windward East Maui. The fog drip to rainfall ratios were estimated using: 1) the fog drip zone boundaries for East Maui (Giambelluca and Nullet, 1991); and 2) an illustration that shows the relationship between fog drip and rainfall for the windward slopes of Mauna Loa, island of Hawaii (Juvik and Nullet, 1995). This method was used to determine the contribution of fog drip in the Kahoma hydrologic unit, which is calculated by multiplying the ratios with the monthly rainfall values in the fog drip zone based on Giambelluca et al (2013). Calculations show that approximately 56 percent of Kahoma lies in the fog drip zone based on elevations greater than 2000 ft. The total contribution from fog drip to the water budget based on percent of fog drig from monthly rainfall is about 33 percent, assuming the same ratios apply to leeward slopes (Table 2-3).

Table 2-3. Fog drip to rainfall ratios for the windward slopes of Mauna Loa on the island of Hawaii and approximate contributions to the Kahoma Hydrologic Unit based on an elevation range of 2000-5780 ft.

Month	Ratio (%)	Mean Rainfall (in)	Contribution (in)
January	13	8.35	1.09
February	13	6.28	0.82
March	13	8.90	1.16
April	27	8.40	2.27
May	27	5.14	1.39
June	27	4.21	1.14
July	67	5.32	3.56
August	67	6.44	4.31
September	67	4.31	2.89
October	40	5.38	2.15
November	40	7.80	3.12
December	27	7.92	2.14

Solar Radiation

Solar radiation is the sun’s energy that arrives at the Earth’s surface after considerable amounts have been absorbed by water vapor and gases in the Earth’s atmosphere. The amount of solar radiation to reach the surface in a given area is dependent in part upon latitude and the sun’s declination angle (angle from the sun to the equator), which is a function of the time of year. Hawaii’s trade winds and the temperature inversion layer greatly affect solar radiation levels, the primary heat source for evaporation. High mountain ranges block moist trade-wind air flow and keep moisture beneath the inversion layer (Lau and Mink, 2006). As a result, windward slopes tend to be shaded by clouds and protected from solar radiation, while dry leeward areas receive a greater amount of solar radiation and thus have higher levels of evaporation. In Kahoma, average annual solar radiation ranged from 129.9 to 296.5 W/m² per day (Figure 2-5). It is greatest at the coast and decreases toward the uplands, where cloud cover is more of an influence (Giambelluca et al. 2014).

Evaporation

Evaporation is the loss of water to the atmosphere from soil surfaces and open water bodies (e.g. streams and lakes). Evaporation from plant surfaces (e.g. leaves, stems, flowers) is termed transpiration. Together, these two processes are commonly referred to as evapotranspiration, and it can significantly affect water yield because it determines the amount of rainfall lost to the atmosphere. On a global scale, the amount of water that evaporates is about the same as the amount of water that falls on Earth as precipitation. However, more water evaporates from the ocean whereas on land, rainfall often exceeds evaporation. The rate of evaporation is dependent on many climatic factors including solar radiation, albedo², rainfall, humidity, wind speed, surface temperature, and sensible heat advection³. Higher evaporation rates are generally associated with greater net radiation, high wind speed and surface temperature, and lower humidity.

² Albedo is the proportion of solar radiation that is reflected from the Earth, clouds, and atmosphere without heating the receiving surface.

³ Sensible heat advection refers to the transfer of heat energy that causes the rise and fall in the air temperature.

A common approach to estimating evaporation is to employ a relationship between potential evaporation and the available water in the watershed. Potential evaporation is the maximum rate of evaporation if water is not a limiting factor, and it is often measured with evaporation pans. In Hawaii, pan evaporation measurements were generally made in the lower elevations of the drier leeward slopes where sugarcane was grown. These data have been compiled and mapped by Ekern and Chang (1985). Most of the drainage basins in Hawaii are characterized by a relatively large portion of the rainfall leaving the basin as evaporation and the rest as streamflow (Ekern and Chang, 1985). Based on the available pan evaporation data for Hawaii, evaporation generally decreases with increasing elevation below the temperature inversion⁴ and the cloud layer (Figure 2-6). At low elevations near the coast, pan evaporation rates are influenced by sensible heat advection from the ocean (Nullet, 1987). Pan evaporation rates are enhanced in the winter by positive heat advection from the ocean, and the opposite occurs in the summer when pan evaporation rates are diminished by negative heat advection (Giambelluca and Nullet, 1992). With increasing distance from the windward coasts, positive heat advection from dry land surfaces becomes an important factor in determining the evaporative demand on the landscape (Nullet, 1987). Shade (1999, Fig. 9) estimated pan evaporation rates of 30 inches per year below 2,000 feet elevation to 90 inches per year near the coast. Within the cloud layer, evaporation rates are particularly low due to the low radiation and high humidity caused by fog drip. Pan evaporation rates dropped below 30 inches per year in this area as reported in Shade (1999, Fig. 9). Near the average height of the temperature inversion, evaporation rates are highly variable as they are mainly influenced by the movement of dry air from above and moist air from below (Nullet and Giambelluca, 1990). Above the inversion, clear sky and high solar radiation at the summit causes increased evaporation, with pan evaporation rates of about 50 to 70 inches per year (Shade, 1999, Fig. 9). For example, Ekern and Chang (1985) reported evaporation increased to 50 percent more than surface oceanic rates near the Mauna Kea crest on the island of Hawaii. Potential annual evapotranspiration in the Kahoma hydrologic unit (Figure 2-6) averages 99.3 inches per year and ranges from 56.0 to 227.8 inches per year (Giambelluca et al. 2014).

Land Use

The Hawaii Land Use Commission (LUC) was established under the State Land Use Law (Chapter 205, Hawaii Revised Statutes) enacted in 1961. Prior to the LUC, the development of scattered subdivisions resulted in the loss of prime agricultural land that was being converted for residential use, while creating problems for public services trying to meet the demands of dispersed communities. The purpose of the law and the LUC is to preserve and protect Hawaii's lands while ensuring that lands are used for the purposes they are best suited. Land use is classified into four broad categories: 1) agricultural; 2) conservation; 3) rural; and 4) urban.

Land use classification is an important component of examining the benefits of protecting instream uses and the appropriateness of surface water use for noninstream uses. While some may argue that land use, in general, should be based upon the availability of surface and ground water resources, land use classification continues to serve as a valuable tool for long-range planning purposes.

As of 2014, the LUC designated 50.2 percent of the land in Kahoma as conservation, 21.9 percent as agriculture, and 27.9 percent as urban (State of Hawaii, Office of Planning, 2015d). The conservation district is located in the upper part of the hydrologic unit, whereas the agricultural district lies in the middle and the urban district is in the lowest part of the hydrologic unit (Figure 2-7).

⁴ Temperature inversion is when temperature increases with elevation.

Land Cover

Land cover for the hydrologic unit of Kahoma is represented by two separate 30-meter Landsat satellite datasets. One of the datasets, developed by the Coastal Change Analysis Program (C-CAP), provides a general overview of the land cover types in Kahoma, e.g., forest, grassland, shrub land, with minor developed areas, cultivated areas, and bare land (Table 2-4, Figure 2-8). The second is developed by the Hawaii Gap Analysis Program (HI-GAP), which mapped the National Vegetation Classification System (NVCS) associations for each type of vegetation, creating a more comprehensive land cover dataset (Table 2-5, Figure 2-7).

Based on the two land cover classification systems, the land cover of Kahoma consists mainly of scrub and forested areas. Over half of the hydrologic unit is made up of alien forest, grassland or shrubland. There is some native dry cliff vegetation, shrubland and sparse ohia spread throughout the upper slopes. A mixture of alien and ohia forests and uluhe shrub lands can be found at intermediate slopes. Alien grasslands and cultivated cropland cover a majority of the lower altitudes near the coast with urban or industrial developments.

Table 2-4. C-CAP land cover classes and area distribution in Kahoma hydrologic unit, Maui, Hawaii. (Source: National Oceanographic and Atmospheric Agency, 2015)

Land Cover	Description	Area (mi ²)	Percent of Unit
Evergreen Forest	Areas where more than 67% of the trees remain green throughout the year	2.341	40.4
Grassland	Natural and managed herbaceous cover	1.241	21.4
Scrub/Shrub	Areas dominated by woody vegetation less than 6 meters in height	1.095	18.9
Cultivated Crops	Areas intensely managed for the production of annual crops	0.247	4.3
Low Intensity Developed	Constructed surface with substantial amounts of vegetated surface	0.230	4.0
Medium Intensity Developed	Areas with a mixture of constructed materials and substantial amounts of vegetation	0.206	3.6
Bare Land	Bare soil, gravel, or other earthen material with little or no vegetation	0.200	3.5
High Intensity Developed	Contains significant land area covered by concrete, asphalt and other constructed materials with less than 20% vegetation	0.138	2.4
Developed Open Space	Areas mostly managed grasses or low-lying vegetation planted for recreation, erosion control, or aesthetic purposes	0.044	0.8
Open Water		0.040	0.7
Palustrine Scrub/Shrub Wetland	Includes tidal and nontidal wetlands dominated by woody vegetation less than 5 meters in height, and all such wetlands that occur in tidal areas in which salinity is below 0.5%	0.005	0.1
Palustrine Emergent Wetland	Includes tidal and nontidal wetlands dominated by persistent emergent vascular plants, mosses or lichens	0.004	0.1

The land cover maps (Figures 2-6, 2-9) provide a general representation of the land cover types in Kahoma. Given that the scale of the maps is relatively large, they may not capture the smaller cultivated lands or other vegetation occupying smaller parcels of land. Land cover types may also have changed slightly since the year when the maps were published. Community members have reported lands cultivated with tropical flowers, dryland (and some wetland) taro, sweet potato, banana, and papaya, long beans, and eggplant primarily in the lower-middle portions of the hydrologic unit. Along the coast, a number of typical native and alien species have been reported. Some of those native species include ulei, naupaka kahakai, akia, lau'we, and hala.

Table 2-5. HI-GAP land cover classes and area distribution in Kahoma hydrologic unit, Maui, Hawaii. (Source: USGS, 2011)

Land Cover	Area (mi ²)	Percent of Unit
Cultivated Cropland	1.890	32.64%
Ohia Forest	0.817	14.12%
Alien Forest	0.675	11.67%
Closed Ohia Forest	0.405	6.99%
Developed, Low Intensity	0.346	5.98%
Alien Shrubland	0.335	5.79%
Native Wet Cliff Vegetation	0.333	5.76%
Alien Grassland	0.277	4.79%
Open Ohia Forest	0.207	3.58%
Uncharacterized Forest	0.141	2.43%
Developed, High Intensity	0.105	1.82%
Uncharacterized Shrubland	0.081	1.39%
Bog Vegetation	0.052	0.90%
Uluhe Shrubland	0.036	0.62%
Uncharacterized Open-Sparse Vegetation	0.035	0.61%
Very Sparse Vegetation to Unvegetated	0.023	0.40%
Native Dry Cliff Vegetation	0.017	0.29%
Native Shrubland / Sparse Ohia (native shrubs)	0.005	0.09%
Open Water	0.005	0.08%
Undefined	0.002	0.04%

Flood

Floods usually occur following prolonged or heavy rainfall associated with tropical storms or hurricanes. The magnitude of a flood depends on topography, ground cover, and soil conditions. Rain falling on areas with steep slopes and soil saturated from previous rainfall events tends to produce severe floods in low-lying areas. Four types of floods exist in Hawaii. Stream or river flooding occurs when the water level in a stream rises into the flood plain. A 100-year flood refers to the probability of a given magnitude flood occurring once in a hundred years, or 1 percent chance of happening in a given year. Flash floods occur within a few hours after a rainfall event, or they can be caused by breaching of a flood safety structure such as a dam. Flash flooding is common in Hawaii because the small drainage basins often have a short response time, typically less than an hour, from peak rainfall to peak streamflow. They are powerful and dangerous in that they can develop quickly and carry rocks, mud, and all the debris in their path down to the coast, causing water quality problems in the near-shore waters. Some floods can even trigger massive landslides, blocking off the entire stream channel. Sheet flooding occurs when runoff builds up on previously saturated ground, flowing from the high mountain slopes to the sea in a shallow sheet (Pacific Disaster Center, 2007). Coastal flooding is the inundation of coastal land areas from excessive sea level rise associated with strong winds or a tsunami.

Peak floods in Kahoma are currently monitored by USGS (1960-present) at USGS station 16638500. The 2-, 5-, 10-, 50-, and 100-year flood magnitudes in Kahoma at this station are calculated as 848, 1670, 2350, 4140, and 5030 cfs. The Federal Emergency Management Agency (FEMA) developed maps that identify the flood-risk areas in an effort to mitigate life and property losses associated with flooding events. Based on these maps, FEMA identified most of the Kahoma hydrologic unit as flood-risk zone X, outside of the the 1% annual chance of flood. Some of the lands along the Kanaha and Kahoma stream channels are designated flood zone A, with a 1% chance of flooding annually. There is a small region

along the coastline in Kahoma where the flood risk is considered zone AE or VE, with the potential for shallow flooding of 1-3 feet annually or areas (Figure 2-10).

Drought

Drought is generally defined as a shortage of water supply that usually results from lower than normal rainfall over an extended period of time, though it can also result from human activities that increase water demand (Giambelluca et al., 1991). The National Drought Mitigation Center (State of Hawaii, Commission on Water Resource Management, 2005b) uses two types of drought definitions — conceptual and operational. Conceptual definitions help people understand the general concept of drought. Operational definitions describe the onset and severity of a drought, and they are helpful in planning for drought mitigation efforts. The four operational definitions of drought are meteorological, agricultural, hydrological, and socioeconomic. Meteorological drought describes the departure of rainfall from normal based on meteorological measurements and understanding of the regional climatology. Agricultural drought occurs when not enough water is available to meet the water demands of a crop. Hydrological drought refers to declining surface and ground water levels. Lastly, socioeconomic drought occurs when water shortage affects the general public.

Impacts of drought are complex and can be categorized into three sectors: water supply; agriculture and commerce; and environment, public health, and safety sectors (State of Hawaii, Commission on Water Resource Management, 2005b). The water supply sector encompasses urban and rural drinking water systems that are affected when a drought depletes ground water supplies due to reduced recharge from rainfall. The agriculture and commerce sector includes the reduction of crop yield and livestock sizes due to insufficient water supply for crop irrigation and maintenance of ground cover for grazing. The environmental, public health, and safety sector focuses on wildfires that are both detrimental to the forest ecosystem and hazardous to the public. It also includes the impact of desiccating streams, such as the reduction of instream habitats for native species.

Droughts have affected the islands throughout Hawaii’s recorded history. The most severe events of the recent past years are associated with the El Niño phenomenon. In January 1998, the National Weather Service’s network of 73 rain gauges throughout the State did not record a single above-normal rainfall, with 36 rain gauges recording less than 25 percent of normal rainfall (State of Hawaii, Commission on Water Resource Management, 2005b). The most recent drought occurred in 2000-2002, affecting all islands, especially the southeastern end of the State.

Table 2-6. Drought risk areas for Maui. (Source: University of Hawaii, 2003)
[Drought classifications of moderate, severe, and extreme have SPI values -1.00 to -1.49, -1.50 to -1.99, and -2.00 or less, respectively]

Sector	Drought Classification (based on 12-month SPI)		
	Moderate	Severe	Extreme
Water Supply	Kula, Kahului, Wailuku, Hana, Lahaina	Kula, Hana	Kula
Agriculture and Commerce	--	--	--
Environment, Public Health and Safety	Kula	Kula	Kula

With Hawaii’s limited water resources and growing water demands, droughts will continue to adversely affect the environment, economy, and the residents of the State. Aggressive planning is necessary to make wise decisions regarding the allocation of water at the present time, and conserving water resources for generations to come. The Hawaii Drought Plan was established in 2000 in an effort to mitigate the long-term effects of drought. One of the projects that supplemented the plan was a drought risk and vulnerability assessment of the State, conducted by researchers at the University of Hawaii (2003). In this project, drought risk areas were determined based on rainfall variation in relation to water source, irrigated area, ground water yield, stream density, land form, drainage condition, and land use. Fifteen

years of historical rainfall data were used. The Standardized Precipitation Index (SPI) was used as the drought index because of its ability to assess a range of rainfall conditions in Hawaii. It quantifies rainfall deficit for different time periods, i.e. 3 months and 12 months. Results of the study for Maui are summarized in Table 2-6. Based on the 12-month SPI, the Kula and Hana regions have the greatest risk to drought impact on Maui because of its dependence on surface water sources, limited rainfall, or relatively high drought frequency and high population density. The growing population in the already densely populated area further stresses the water supply. The Kahoma region is vulnerable to drought in the agricultural sector and has substantial wildfire risk, but there is less risk of drought to the general population.

Figure 2-2. Generalized geology of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Sherrod et al., 2007)




Figure 2-3. Soil classification of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015m)




Figure 2-4. Mean annual rainfall of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Giambelluca et al., 2013)




Figure 2-5. Mean annual solar radiation of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015c)




Figure 2-6. Mean annual potential evapotranspiration (Penman-Monteith method) of the Kahoma hydrologic unit, Maui, Hawaii. (Source: Giambelluca et al., 2014)




Figure 2-7. State land use district boundaries of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015d).




Figure 2-8. C-CAP land cover of the Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Office of Planning, 2015k).




Figure 2-9. Hawaii GAP land cover classes of the Kahoma hydrologic unit, Maui, Hawaii (Source: USGS, 2011).





Figure 2-10. FEMA flood zone regions in Kahoma hydrologic unit (Source: Federal Emergency Management Agency, 2014).



3.0 Hydrology

The Commission, under the State Water Code, is tasked with establishing instream flow standards by weighing “the importance of the present or potential instream values with the importance of the present or potential uses of water for noninstream purposes, including the economic impact of restricting such uses.” While the Code outlines the instream and offstream uses to be weighed, it assumes that hydrological conditions will also be weighed as part of this equation. The complexity lies in the variability of local surface water conditions that are dependent upon a wide range of factors, including rainfall, geology, and human impacts, as well as the availability of such information. The following is a summary of general hydrology and specific hydrologic characteristics for Kahoma Stream.

Streams in Hawaii

Streamflow consists of: 1) direct surface runoff in the form of overland flow and subsurface flow that rapidly returns infiltrated water to the stream; 2) ground water discharge in the form of base flow; 3) water returned from streambank storage; 4) rain that falls directly on streams; and 5) additional water, including excess irrigation water discharged into streams by humans (Oki, 2003). The amount of runoff and ground water that contribute to total streamflow is dependent on the different components of the hydrologic cycle, as well as man-made structures such as diversions and other stream channel alterations (e.g. channelizations and dams).

Streams in Hawaii can either gain or lose water at different locations depending on the geohydrologic conditions. A stream gains water when the ground water table is above the streambed. When the water table is below the streambed, the stream can lose water. Where the streambed is lined with concrete or other low-permeability or impermeable material, interaction between surface water and ground water is unlikely. Another way that ground water influences streamflow is through springs. A spring is formed when a geologic structure (e.g., fault or fracture) or a topographic feature (e.g., side of a hill or a valley) intersects ground water either at or below the water table. It can discharge ground water onto the land surface, directly into the stream, or into the ocean. Figure 3-1 illustrates a valley that has been incised into a high-level water table, resulting in ground water discharges that contribute directly to streamflow and springs that contribute to streamflow.

The USGS has maintained a variety of stream and ditch gaging stations over time, although none are currently active (Figure 3-2). These stations have improved our understanding of flood frequency and intensity (station 16638500), the flow of water in the Lahainaluna Ditch (station 16637000), the the flow of water in the headwaters (stations 16636000, 16634000) or from development tunnels (station 16633400). Historic data from these stations are available in Table 3-1.

Figure 3-1. Conceptual diagram illustrating surface water-groundwater interactions in West Maui. (Source: Oki et al., 2010).




Table 3-1. Selected discharge exceedance values from historic stream gaging data in the Kahoma hydrologic unit, Maui, Hawaii. (Source: USGS NWIS, 2018). [in cubic feet per second (million gallons per day)]

Station	Stream	Location	Period of record	Discharge (Q) for a selected percentage (xx) discharge was equaled or exceeded				
				Q ₅₀	Q ₆₀	Q ₇₀	Q ₈₀	Q ₉₀
16634000	Kahoma	above diversion	1911-1917	5.7 (3.68)	4.2 (2.71)	3.9 (2.52)	0.31 (0.20)	0.0 (0.0)
16636000	Kanaha	above diversion	1911-1917	4.9 (3.17)	4.5 (2.91)	4.3 (2.78)	3.7 (2.39)	3.4 (2.20)
16638000	Kanaha	below diversion above confluence with Kahoma	1911-1916	1.6 (1.03)	1.4 (0.90)	1.2 (0.78)	1.0 (0.65)	0.77 (0.50)
16638500	Kahoma	below confluence with Kanaha	1963-1990	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

Ground Water

Ground water is an important component of streamflow as it constitutes the base flow¹ of Hawaiian streams. When ground water is withdrawn from a well, the water level in the surrounding area is lowered. Nearby wetlands or ponds may shrink or even dry up if the pumping rate is sufficiently high (Gingerich and Oki, 2000). The long-term effects of ground water withdrawal can include the reduction of streamflow, which may cause a decrease in stream habitats for native species and a reduction in the amount of water available for irrigation. The interaction between surface water and ground water warrants a close look at the ground water recharge and demand within the State as well as the individual hydrologic units.

In Hawaii, ground water is replenished by recharge from rainfall, fog drip, and irrigation water that percolate through the plant root zone to the subsurface rock. Recharge can be captured in three major fresh ground water systems: 1) fresh water-lens system; 2) dike-impounded system; and 3) perched system. The fresh water-lens system provides the most important sources of ground water. It includes a lens-shaped layer of fresh water, an intermediate transition zone of brackish water, and underlying salt water. Vertically extensive fresh water-lens system can extend several hundreds or even thousands of feet below mean sea level. By contrast, a dike-impounded system is found in rift zones and caldera of a volcano where low-permeability dikes compartmentalize areas of permeable volcanic rocks, forming high-level water bodies. On Maui, dikes impound water to as high as 3,300 feet above mean sea level. A perched system is found in areas where low-permeability rocks impede the downward movement of percolated water sufficiently to allow a water body to form in the unsaturated zone above the lowest water table (Gingerich and Oki, 2000).

The hydrologic unit of Kahoma spans the Launiupoko and Honokowai aquifer systems. A general overview of the ground water occurrence, movement, and interactions with surface water in this area is described in Oki (2010) and illustrated in Figure 3-1. Ground water is found in dike-impounded structures of Honolua and Wailuku Volcanics as well as in the fresh water-lens system making up the basal aquifer. Withdrawal from wells at or below sea level should not affect the high-elevation water table because the thick unsaturated zone will prevent any significant changes in the vertical flow gradient.

¹ Base flow is the water that enters a stream from persistent, slowly varying sources (such as the seepage of ground water), and maintains stream flow between water-input events (i.e., it is the flow that remains in a stream in times of little or no rainfall).

However, during certain hydrologic conditions, there may be surface water-groundwater interactions, especially in gaining reaches.

Wells in the Kahoma Hydrologic Unit

The Lahaina aquifer sector includes many aquifer systems: Honokohau, Honolua, Honokowai, Launiupoko, Olowalu, and Ukumehame. Historic records put some perspective on the current pumping rates from the Lahaina aquifer sector. Groundwater withdrawals steadily increased with the expansion of plantation agriculture and urbanization. From 1918-1929, an average of 33 mgd was withdrawn from the aquifer, while from 1930-1979, withdrawals averaged 44 mgd, primarily from the Honokowai and Launiupoko aquifer systems (Gingerich and Engott, 2012). Following growth of tourism and eventual closure of pineapple and sugar cane plantation operations, groundwater withdrawals declined to about 21 mgd from 1980 to 2000, and eventually down to 4 mgd by 2008. However, the 2017 12-month moving average for the Lahaina aquifer sector was 7.148 mgd, demonstrating an increased demand by tourism and a growing resident population (State of Hawaii, Commission on Water Resource Management 2018b).


In 2017, the 12-month moving average for existing groundwater withdrawals from the Launiupoko aquifer system was 0.864 mgd and for the Honokowai aquifer system was 3.568 mgd. This is below each of aquifer sector's current sustainable yield of 7 mgd and 6 mgd, respectively (State of Hawaii, Commission on Water Resource Management, 2015d). This water is to meet municipal needs in the Lahaina and Kaanapali areas. Detailed information for each well is specified in Table 3-2. Total monthly groundwater pumpage and the 12-month moving average from 1990 to 2018 are provided in Figure 3-3 for the Launiupoko aquifer sector. The location of wells in the hydrologic unit in relation to the aquifer sectors in Lahaina are depicted in Figure 3-2.

Nearby, Pioneer Mill drilled two shaft wells (5240-002 & 5240-003) for sugar cane irrigation in 1897 to skim freshwater off of the basal aquifer. These shafts are largely no longer in use, although a small pump was located in well 5240-003 (CWRM site visit Sept 2018). They also built three high elevation tunnels to increase the supply of dike-impounded groundwater to streamflow. Two of these three tunnels are collapsed (5436-001 & 5437-001) and are abandoned, with the third (5437-002), and most productive of the tunnels, continuing to contribute to surface flows above the main intake on Kahoma Stream. Maui DWS (6-5339-003 & 6-5339-003) and Maui Parks and Recreation have drilled wells for water supply in the Kahoma hydrologic unit. A majority of the reported ground water use is for municipal use and irrigation of landscaping, with a small amount used for agriculture (Table 3-2).

Groundwater Pumping and Salinity Levels

Of growing concern is the impact of increased groundwater pumpage on groundwater salinity. As pumpage draws down the less dense freshwater lense, brackish groundwater from the transition zone will mix, increasing the salinity (and thus chloride content) of the freshwater lense. Freshwater is water with a chloride content of less than 250 parts per million (ppm), or 1.3% of the chloride content of seawater. Hydrologists with the USGS and CWRM monitor the elevation of the freshwater lense, the transition zone to brackish water, and the elevation of sea water in deep monitoring wells across the state. In West Maui, the Mahinahina Deep Monitoring Well has been an important tool for monitoring groundwater conditions in the Honokowai Aquifer sector for many years (Figure 3-4). There is no deep monitoring well for the Launiupoko Aquifer sector. In areas of the aquifer that have a thin freshwater lense, the impact of over pumping of groundwater on salinity levels can be immediately apparent (Figure 3-5). Thus, even if current pumpage does not reach the aquifer's sustainable yield, pumpage can negatively impact the future use of the aquifer.

Figure 3-2. Well locations and numbers in the Kahoma hydrologic unit, which spans the Launiupoko and Honokowai aquifer sectors, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b).



Prepared by the Department of Land and Natural Resources,
 Commission on Water Resource Management
 Transverse Mercator Projection, Zone 4, North American Datum 1983

Table 3-2. Information of wells located in Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b). [IRRLA = Landscape irrigation; MUNCO = Municipal, County; DOM = domestic; UNU = unused; AGRCP = Agriculture row crop; HHFDC = Hawaii Housing Finance & Development Corporation]
 [Negative elevation values indicate feet below mean sea level; positive elevation values indicate feet above mean sea level. Pump rate measured in gallons per minute (gpm); n/a indicates value is unknown.]

Well number	Well Name	Well Owner	Year drilled	Use	Ground elevation (feet)	Well depth (feet)	Pump elevation (feet)	Pump depth (feet)	Pump rate (mgd)	2017 12-MAV (mgd)
6-5240-001	Mill Shaft Pump C	Pioneer Mill	1897	ABN	34	39	--	--	--	n/a
6-5240-009	Kahoma Irrigation	Opukea at Lahaina	2006	IRRLA	29	58	-8	37	0.216	--
6-5241-001	Lahaina Surf	Hale Mahaolu	1985	IRRLA		28			0.050	0.004
6-5339-001	Waipuka 1	Maui County	1962	MUNCO	440	482	-18	458	0.324	0.000
6-5339-002	Waipuka 2	Maui County	1963	MUNCO	441	498	-26.65	467.65	0.360	0.095
6-5339-003	Kanaha 1	Maui County	1977	MUNCO	590	642	-14.75	604.75	0.360	0.059
6-5339-004	Kanaha 2	Maui County	1974	MUNCO	654	749	-21	675	0.360	0.044
6-5340-001	Kahoma Pump M	Kahoma Land LLC	1933	AGRCP	322	323			10.08	0.000
6-5341-002	Front St House	William Dorbush	2000	IRRLA	13	35	-16	29	0.147	--
6-5436-001	Kahoma Tun 3	Kahoma Ranch	--	n/a	2,350	n/a	n/a	n/a	n/a	--
6-5437-001	Kahoma Tun 1	Kahoma Ranch	--	n/a	1,923	n/a	n/a	n/a	n/a	--
6-5437-002	Kahoma Tun 2	Kahoma Ranch	--	AGRCP	1,984	n/a	n/a	n/a	n/a	1.67
6-5439-001	Wahikuli 1	HHFDC	1992	DOM	1,055	1,115	n/a	n/a	n/a	0.000
6-5439-002	Wahikuli 2	HHFDC	1993	UNU	1,055	1,120	n/a	n/a	n/a	0.000
6-5440-001	Wahikuli Irrigation	HHFDC	1993	IRRLA	148	162	n/a	n/a	n/a	0.017
6-5441-002	Lahaina Civic Center	Maui County	1993	IRRLA	n/a	97	n/a	n/a	n/a	0.148

Figure 3-3. Monthly reported pumpage, 12-month moving average (12-MAV) and sustainable yield for the Launiupoko Aquifer sector. (Source: Source: State of Hawaii, Commission on Water Resource Management, 2018b).





Figure 3-4. Elevation to freshwater lens, brackish water transition and sea water in the Mahinahina Deep Monitoring Well, West Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018a).



Notes: (1) TTZ = 1,000 $\mu\text{S/cm}$ (~ 220 mg/L Cl^-); MPTZ = 25,000 $\mu\text{S/cm}$ (~ 8,500 mg/L Cl^-) (2) Fresh Water < 220 mg/L Cl^- , Brackish Water 220 mg/L Cl^- to 16,999 mg/L Cl^- , Sea Water \geq 17,000 mg/L Cl^- ; (3) OS 421/425 = Ocean Sensors CTD (absolute conductivity); (4) RBR 12895 = RBR Global CTD (Specific Conductivity); (5) msl = mean sea level. Conditions inside the well prevented successful CTD deployment from 2006 through 2012. CTD profiling of this well was resumed 3-13-2013.

* Since the year 2001, the MPTZ rose 16.11 feet, to an elevation of 123.63 feet below mean sea level, where it is near the calculated Ghyben-Herzberg equilibrium elevation of 136 feet below msl, relative to the Water Table, measured at 3.40 feet above msl.

Kahoma and Kanaha Streamflow Characteristics

Dike-impounded groundwater supports continuous flow in the upper reaches of Kahoma and Kanaha streams while the lowest reaches are losing reaches, with streamflow contributing to groundwater recharge (Figure 3-6). Stream flow above the diversion on Kahoma and Kanaha streams (natural flow) was monitored by the U.S. Geological Survey (USGS) from 1911 to 1917 (station #16634000) and from 1911 to 1916 (station # 16636000), respectively (Figure 3-6). CWRM has daily flow data for Kahoma Ditch and Lahainaluna Ditch from 1989 to 1994 (Table 14-3, Figure 14-3). The USGS established partial-record gaging stations on Kahoma (station 205404156372401) and Kanaha (station 205334156382201) as part of a low-flow study in West Maui (Cheng 2014) to determine low-flow exceedance values by record extension using long-term index stations. The closest long-term continuous stream flow monitoring stations maintained by the USGS are on Honokohau Stream (station #16620000) and Wailuku River (station #16604500), but Halawa Stream on Mokokai (station #16400000) was chosen as the index station for Kanaha Stream’s record augmentation in Cheng (2014). From 2012 to 2013, a low-flow partial record gaging station was established at a station upstream of the uppermost diversion on Kahoma and Kanaha streams and natural-flow duration discharges were estimated for the current period (1984-2013) as provided in Table 3-3.

Figure 3-5. Monthly groundwater pumpage and chloride concentration from two municipal (A & B) and one private (C) well in West Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018b).




Table 3-3. Selected natural-flow duration discharge exceedance values for the current (1984-2013) climate period based on partial record gaging station for the Kahoma hydrologic unit, Maui, Hawaii. (Source: Cheng 2014) [Flows are in cubic feet per second (million gallons per day)]

Stream	Discharge (Q) for a selected percentage (xx) discharge was equaled or exceeded				
	Q ₅₀	Q ₆₀	Q ₇₀	Q ₈₀	Q ₉₀
Kahoma	5.8 (3.75)	5.6 (3.62)	5.4 (3.49)	4.9 (3.17)	2.9 (1.87)
Kanaha	4.9 (3.17)	4.7 (3.04)	4.5 (2.91)	4.3 (2.78)	4.1 (2.65)

Kahoma and Kanaha streams are gaining in the uppermost reaches in the watershed, but in the reach immediately below the main diversions on each stream, from about 1850 ft in elevation in Kahoma and 1250 ft in Kanaha, to the ocean, the streams have losing reaches (Figure 3-7). Using seepage runs, USGS estimated seepage loss below the upper diversion on Kahoma Stream to be between 0.62 and 1.18 cubic feet per second per mile (0.40 and 0.76 million gallons per day per mile). Similarly, there was an estimated 0.60 cubic feet per second per mile (0.39 million gallons per day per mile) seepage loss on Kanaha Stream. There is approximately 3.40 miles between diversion 951 on Kahoma Stream and the confluence with Kanaha Stream. There is approximately 1.85 miles between diversion 954 on Kanaha Stream and the confluence with Kahoma Stream. The auwai that feeds the loi complex in Kahoma Valley is approximately 2.25 miles below diversion 951. USGS estimated that Kahoma Stream would naturally flow to the confluence with Kanaha Stream at least 85-950% of the time (and recent gaging evidence suggests 100% of the time). Kanaha Stream was estimated to naturally flow to the confluence with Kahoma Stream at least 95% of the time (Cheng 2014). Lost streamflow recharges the groundwater systems.

As part of an effort to improve resource monitor, Kamehamea Schools Bishop Estate (KSBE) installed a real-time continuous monitoring station on Kahoma Stream at an elevation of 550 feet. The primary purpose was to determine stream depth (and therefore how much water was available) over time. While these data provide some qualitative data, there have been issues with the site location due to: (1) changing stream channel conditions, the control point has shifted over time; (2) discharge measurements were not consistently made to develop an accurate rating curve to convert depth of stage to streamflow. In 2017, in cooperation with KSBE, CWRM staff installed a staff plate for reference and began measuring discharge consistently. These measurements are presented in Table 3-4, along with an estimate of the flow rate of water withdrawn at diversion 951 and the estimated total flow natural flow at the station.

Table 3-4. Discharge measurements at gage 6-123 and estimated diversion amounts from diversion 951 on Kahoma Stream. (Source: CWRM) [Flows are in cubic feet per second (million gallons per day)]

date	time	discharge	diverted amount	estimated natural flow
10/2/2017	10:15	0.87 (0.56)	0	0.87 (0.56)
10/27/2017	14:13	1.56 (1.01)	0.23 (0.15)*	1.79 (1.16)
11/30/2017	10:20	23.16 (14.97)	0.23 (0.15)*	23.39 (15.12)
12/12/2017	15:25	1.02 (0.66)	0	1.02 (0.66)
12/28/2017	12:04	1.29 (0.84)	0	1.29 (0.84)
03/06/2018	09:55	1.73 (1.12)	0.23 (0.15)*	1.96 (1.27)
08/14/2018	09:12	2.87 (1.86)	0.23 (0.15)*	3.10 (2.01)

*estimate

Long-term trends in flow

In a different study, the USGS examined the long-term trends and variations in streamflow on the islands of Hawaii, Maui, Molokai, Oahu, and Kauai, where long-term stream gaging stations exist (Oki, 2004). The study analyzed both total flow and estimated base flow at 16 long-term gaging stations. Figure 3-5 illustrates the results of the study for 7 long-term gaging stations around the islands. According to the analyses, low flows generally decreased from 1913 to 2002, which is consistent with the long-term downward trends in rainfall observed throughout the islands during that period. Monthly mean base flows decreased from the early 1940s to 2002, which is consistent with the measured downward trend of low flows from 1913 to 2002. This long-term downward trend in base flow may imply a reduction of ground water contribution to streams. Kowaikoi streamflow declined from the 1920s to present day while Waiale streamflow has increased during this same period, suggesting a shift in regional rainfall patterns or changes in the hydrology of the Alakai Marsh. Changing streamflow characteristics could pose a negative effect on the availability of drinking water for human consumption and habitat for native stream fauna (Oki, 2004).

Figure 3-6. USGS surface water gage locations by number for the Kahoma hydrologic Unit, Maui, Hawaii. (Source: USGS, 2011).




Figure 3-7. Seepage run results and median flow in cubic feet per second from partial record gaging stations of the Kahoma hydrologic unit (Source: State of Hawaii, Commission on Water Resource Management, 2015a; Cheng, 2014).





Figure 3-8. Cumulative departures of monthly mean flow from the mean of the monthly flows, Hawaii. This data is based on complete water years from 1913 through 2002. (Oki, 2004, Figure 4)



4.0 Maintenance of Fish and Wildlife Habitat


When people in Hawaii consider the protection of instream flows for the maintenance of fish habitat, their thoughts generally focus on just a handful of native species including five native fishes (four gobies and one eleotrid), two snails, one shrimp, and one prawn. Table 4-1 below identifies commonly mentioned native stream animals of Hawaii.

Table 4-1. List of commonly mentioned native stream organisms. (Source: State of Hawaii, Division of Aquatic Resources, 1993)

Scientific Name	Hawaiian Name	Type
<i>Awaous stamineus</i>	‘O‘opu nakea	Goby
<i>Lentipes concolor</i>	‘O‘opu hi‘ukole (alamo‘o)	Goby
<i>Sicyopterus stimpsoni</i>	‘O‘opu nopili	Goby
<i>Stenogobius hawaiiensis</i>	‘O‘opu naniha	Goby
<i>Eleotris sandwicensis</i>	‘O‘opu akupa (okuhe)	Eleotrid
<i>Atyoida bisulcata</i>	‘Opae kala‘ole	Shrimp
<i>Macrobrachium grandimanus</i>	‘Opae ‘oeha‘a	Prawn
<i>Neritina granosa</i>	Hihiwai	Snail
<i>Neritina vespertina</i>	Hapawai	Snail

Hawaii’s native stream animals have amphidromous life cycles (Ego, 1956) meaning that they spend their larval stages in the ocean (salt water), then return to freshwater streams to spend their adult stage and reproduce. Newly hatched fish larvae are carried downstream to the ocean where they become part of the planktonic pool in the open ocean. The larvae remain at sea from a few weeks to a few months, eventually migrating back into a fresh water stream as juvenile *hinana*, or postlarvae (Radtke et al., 1988). Once back in the stream, the distribution of the five native fish species are largely dictated by their climbing ability (Nishimoto and Kuamoo, 1991) along the stream’s longitudinal gradient. This ability to climb is made possible by a fused pelvic fin which forms a suction disk. *Eleotris sandwicensis* lacks fused pelvic fins and is mostly found in lower stream reaches. *Stenogobius hawaiiensis* has fused pelvic fins, but lacks the musculature necessary for climbing (Nishimoto and Kuamoo, 1997). *Awaous stamineus* and *Sicyopterus stimpsoni* are able to ascend moderately high waterfalls (less than ~20 meters) (Fitzsimons and Nishimoto, 1990), while *Lentipes concolor* has the greatest climbing ability and has been observed at elevations higher than 3,000 feet (Fitzsimons and Nishimoto, 1990) and above waterfalls more than 900 feet in vertical height (Englund and Filbert, 1997). Figure 4-1 illustrates the elevational profile of these native fresh water fishes.

Figure 4-1. Elevational profile of a terminal-estuary stream on the Big Island of Hawaii (Hakalau Stream). (Source: McRae, 2007, adapted from Nishimoto and Kuamoo, 1991 [with permission])



The maintenance, or restoration, of stream habitat requires an understanding of and the relationships among the various components that impact fish and wildlife habitat, and ultimately, the overall viability of a desired set of species. These components include, but are not limited to, species distribution and diversity, species abundance, predation and competition among native species, similar impacts by alien species, obstacles to migration, water quality, and streamflow. The Commission does not intend to delve into the biological complexities of Hawaiian streams, but rather to present basic evidence that conveys the general health of the subject stream. The biological aspects of Hawaii's streams have an extensive history, and there is a wealth of knowledge, which continues to grow and improve.

Hawaii Stream Assessment

One of the earliest statewide stream assessments was undertaken by the Commission in cooperation with the National Park Service's Hawaii Cooperative Park Service Unit. The 1990 Hawaii Stream Assessment (HSA) brought together a wide range of stakeholders to research and evaluate numerous stream-related attributes (e.g., hydrology, diversions, gaging, channelizations, hydroelectric uses, special areas, etc.). The HSA specifically focused on the inventory and assessment of four resource categories: 1) aquatic; 2) riparian; 3) cultural; and 4) recreational. Though no field work was conducted in its preparation, the HSA involved considerable research and analysis of existing studies and reports. The data were evaluated according to predefined criteria and each stream received one of five ranks (outstanding, substantial, moderate, limited, and unknown). Based on the stream rankings, the HSA offered six different approaches to identifying candidate streams for protection: streams with outstanding resources (aquatic, riparian, cultural or recreational), streams with diverse or "blue ribbon" resources, streams with high quality natural resources, streams within aquatic resource districts, free flowing streams, or streams within the National Wild and Scenic Rivers database.

Due to the broad scope of the HSA inventory and assessment, it continues to provide a valuable information base for the Commission's Stream Protection and Management Program and will continue to be referred to in various sections throughout this report. The HSA recommended that the Kahoma hydrologic unit streams not be listed as candidate streams for protection based on its riparian, aquatic, cultural and recreational resources. At the time of the assessment, only one out of four group one native species had been observed in Kahoma Stream: nakea (*Awaous stamineus*). Additionally, only one group two native species had been observed in the stream. While the Kahoma Stream originally provided habitat to a number of native species, due to significant land degradation from non-native species and the introduction of non-native aquatic species, the HSA downgraded the stream to having only moderate aquatic resources.

The HSA inventory was general in nature, resulting in major data gaps, especially those related to the distribution and abundance of aquatic organisms – native and introduced – inhabiting the streams. The State of Hawaii Division of Aquatic Resources (DAR) has since continued to expand the knowledge of aquatic biota in Hawaiian streams. Products from their efforts include the compilation and publication of an *Atlas of Hawaiian Watersheds and Their Aquatic Resources* for each of five major islands in the state (Kauai, Hawaii, Oahu, Molokai, and Maui). Each atlas describes watershed and stream features, distribution and abundance of stream animals and insect species, and stream habitat use and availability. Based on these data, a watershed and biological rating is assigned to each stream to allow easy comparison with other streams on the same island and across the state. The data presented in the atlases are collected from various sources, and much of the stream biota data are from stream surveys conducted by DAR. Currently, efforts have been focused on updating the atlases with more recent stream survey data collected statewide, and developing up-to-date reports for Commission use in interim IFS recommendations. A copy of the updated inventory report for Kahoma Stream is in Appendix A. The following is a brief summary of findings.


- **Point Quadrat Survey.** In Kahoma Stream, only two point quadrat surveys were conducted in 1963, likely due to the lack of stream flow below the diversion and lack of access above the diversion. Although the post larval recruitment of native fish was not surveyed, diversions that fully dewater streams restrict the passage of larval and adult stream animals. Streamflow restoration would likely benefit recruitment and survival of native species in the lower, middle and upper reaches.
- **Insect Survey.** Kahoma Stream did not meet the DAR qualification for native insect diversity (>19 spp) and native macrofauna diversity (>5 spp.) giving it a low score for native species, likely due to a lack of streamflow. Kahoma Stream supports native damselflies (*Megalagrion spp.*) in the middle and upper stream reaches. Some native damselfly are currently proposed for listing as Endangered under the federal Endangered Species Act. Streamflow restoration may increase insect biota diversity.
- **Watershed and Biological Rating.** The Kahoma watershed has an average rating for Maui and an average rating statewide for land cover. The lack of estuarine and shallow marine areas associated with Kahoma ranks it low for Maui and the state, while its stewardship rating is about average for Maui and above average for the state. Kahoma Stream has an above average rating for stream size, a below average rating for wetness, and an above average reach diversity resulting in an above average total watershed rating for Maui and statewide. The below average number and diversity of native species found throughout the watershed, with a high rating of introduced species gives it an average all species and total biological rating for the island and the state. These scores combined gave Kahoma Stream an average overall watershed rating.

A total of two point-quadrat surveys were conducted in 1963, identifying *Atyoida bisulcata* at one site and an unknown gobiid species at another. Tingley et al. (2018) developed a classification of stream reaches that identifies the suitability for native species based on hydrologic, anthropogenic, and landscape factors (Figure 4-2). In 2017, Steward (unpublished) conducted point-quadrat surveys in middle and upper reaches in Kahoma Stream. The dominant native species observed were *Awaous stamineus* with 95 observed across a variety of habitats (Table 4-2).

Table 4-2. Size distribution of *Awaous stamineus* by habitat type in Kahoma Stream.

habitat	total	1-3 inch	3-5 inch	5-7 inch
Pool	20	20	0	0
Riffle	3	1	2	0
Run	56	39	16	1
Side pool	14	14	0	0

Figure 4-2. Classification of stream reaches based on habitat suitability for native species. (Source: Tingley III et al. 2018)



5.0 Outdoor Recreational Opportunities


Water-related recreation is an integral part of life in Hawaii. Though beaches may attract more users, the value of maintaining streamflow is important to sustaining recreational opportunities for residents and tourists alike. Streams are often utilized for water-based activities, such as boating, fishing, and swimming, while offering added value to land-based activities such as camping, hiking, and hunting. Growing attention to environmental issues worldwide has increased awareness of stream and watershed protection and expanded opportunities for the study of nature; however, this must be weighed in conjunction with the growth of the eco-tourism industry and the burdens that are placed on Hawaii's natural resources.

The Hawaii Stream Assessment identified hiking and scenic views as stream recreational opportunities in the Kahoma hydrologic unit and the regional committee classified the stream as "substantial" (National Park Service, Hawaii Cooperative Park Service Unit, 1990). The Kahoma Ranch offers zip-line and all-terrain vehicle (ATV) tours through former agricultural lands. The roads used to access these opportunities are often watered with diverted stream water for dust control.

Mammal hunting is permitted in Unit A on the island of Maui in the state-owned portions of the West Maui Forest Reserve that includes the Kahoma hydrologic unit. None of the Kahoma hydrologic unit is in designated as hunting area (Figure 5-1).

Since changes to streamflow and stream configurations have raised concerns regarding their impact to on-shore and near-shore activities, the Commission attempted to identify these various activities in relation to Kahoma Stream. A 1981 Maui Resource Atlas, prepared by the State of Hawaii Department of Transportation's Harbors Division, inventoried coral reefs and coastal recreational activities. Looking at available GIS data, some of the activities the Commission identified that were known to occur or observed at or near Kahoma include: pole and line fishing, bait fishing, gill netting, torch fishing, sport diving and throw netting, surfing, standup paddle boarding, seaweed and shell collecting (Figure 5-1).

Figure 5-1. Public hunting areas for game mammals and coastal resources in Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 1999, 2002b, 2002c, 2004a)




Prepared by the Department of Land and Natural Resources,
 Commission on Water Resource Management
 Transverse Mercator Projection, Zone 4, North American Datum 1983

6.0 Maintenance of Ecosystems

An ecosystem can be generally defined as the complex interrelationships of living (biotic) organisms and nonliving (abiotic) environmental components functioning as a particular ecological unit. Depending upon consideration of scale, there may be a number of ecosystem types that occur along a given stream such as estuaries, wetlands, and stream vegetation, according to the State Water Code. Figure 6-1 provides a simplified ecosystem represented in a Hawaiian stream. The entire hydrologic unit, as it relates to hydrologic functions of the stream, could also be considered an ecosystem in a very broad context.

The HSA determined that Kahoma Stream did not deserve to be a candidate stream for protection based on its Diversity of Resources scoring (does not have an outstanding rating in three categories) or its Blue Ribbon Resources scoring. Detrimental organisms (non-native species that have a negative effect on the ecosystem) were not considered in the final ranking; however their presence and abundance are considerable ecosystem variables. The destruction to both vegetation and soil resources in the Kahoma hydrologic unit by detrimental organisms (pigs) is considered a major factor in alterations to the hydrology and water quality of the stream.

Figure 6-1. Simplified ecosystem illustrated in a Hawaiian stream. (Source: Ziegler, 2002, illustration by Keith Kruger).



The Hawaiian resource-use concept of ahupuaa is closely related to the Western concepts of ecosystem maintenance. Native Hawaiians generally utilized natural resources within the limits of their ahupuaa; therefore, it was important to manage and conserve these resources. Likewise, watershed resources must be properly managed and conserved to sustain the health of the stream and the instream uses that are dependent upon it.

The riparian resources of Kahoma Stream were classified by the HSA (National Park Service, Hawaii Cooperative Park Service Unit, 1990) and ranked according to a scoring system using six of the seven variables (Table 6-1).

Table 6-1. Hawaii Stream Assessment indicators of riparian resources for Kahoma hydrologic unit, Maui, Hawaii.

Category	Value
<p>Listed threatened and endangered species: These species are generally dependent upon undisturbed habitat. Their presence is, therefore an indication of the integrity of the native vegetation. The presence of these species along a stream course was considered to be a positive attribute; with the more types of threatened and endangered species associated with a stream the higher the value of the resource. Only federally listed threatened or endangered forest or water birds that have been extensively documented within the last 15 years were included.</p>	0
<p>Recovery habitat: Recovery habitat consists of those areas identified by the USFWS and DLNR as essential habitat for the recovery of threatened and endangered species. Streams that have recovery habitat anywhere along their length were included.</p>	none
<p>Other rare organisms and communities: Many species that are candidates for endangered or threatened status have not been processed through all of the requirements of the Endangered Species Act. Also a number of plant communities associated with streams have become extremely rare. These rare organisms and communities were considered to be as indicative of natural Hawaiian biological processes as are listed threatened and endangered species.</p>	4
<p>Protected areas: The riparian resources of streams that pass through natural area reserves, refuges and other protected areas are accorded special protection from degradation. Protected areas were so designated because of features other than their riparian resources. The presence of these areas along a stream, however, indicates that native processes are promoted and alien influences controlled.</p>	Yes
<p>Wetlands: Wetlands are important riparian resources. They provide habitat for many species and are often important nursery areas. Because they are often extensive areas of flat land generally with deep soil, many have been drained and converted to agricultural or urban uses. Those that remain are, therefore, invaluable as well as being indicators of lack of disturbance.</p>	W (Palustrine Wetland)
<p>Native forest: The proportion of a stream course flowing through native forest provides an indication of the potential "naturalness" of the quality of a stream's watershed; the greater the percentage of a stream flowing through native forest most of which is protected in forest reserves the more significant the resource. Only the length of the main course of a stream (to the nearest 10 percent) that passes through native forest was recorded.</p>	50%
<p>Detrimental organisms: Some animals and plants have a negative influence on streams. Wild animals (e.g., pigs, goats, deer) destroy vegetation, open forests, accelerate soil erosion, and contaminate the water with fecal material. Weedy plants can dramatically alter the nature of a stream generally by impeding water flow. Three species, California grass, hau, and red mangrove, are considered to have the greatest influence. The presence of any of these animals or plants along a stream course was considered a potentially negative factor, while the degree of detriment is dependent on the number of species present.</p>	Pigs

For the purpose of this section, management areas are those locales that have been identified by federal, state, county, or private entities as having natural or cultural resources of particular value. The result of various government programs and privately-funded initiatives has been a wide assortment of management areas with often common goals. Such designated areas include forest reserves, private preserves, natural area reserves, wildlife sanctuaries, national parks, historic landmarks, and so on. In Kahoma, about 11.5 percent of the hydrologic unit falls within the West Maui Natural Area Reserve or the West Maui Forest Reserve (Table 6-2).

Table 6-2. Management areas located within Kahoma hydrologic unit, Maui, Hawaii. (Source: State of Hawaii, Division of Forestry and Wildlife, 2008a).

Management Area	Managed by	Area in Unit (mi ²)	Percent of Unit
West Maui Natural Area Reserve Panaewa Section	State Division of Forestry and Wildlife	1.77	30.9
The West Maui Natural Area Reserve is one of 21 natural area reserves across the state managed by the State Department of Land and Natural Resources (DLNR)'s Division of Forestry and Wildlife. The management goal of the Natural Area Reserve System is to preserve and protect representative samples of Hawaiian biological ecosystems and geological formations. Management objectives of natural area reserves are to: 1) control non-native animals; 2) control non-native pest plants; 3) restore and enhance native habitat; 4) monitor the ecosystem; 5) provide public information and education; 6) encourage sound scientific research			
West Maui Natural Area Reserve Honokowai Section	State Division of Forestry and Wildlife	< 0.01	< 0.1
Kapunakea Preserve	The Nature Conservancy	< 0.01	< 0.1
West Maui Forest Reserve	State Division of Forestry and Wildlife	0.015	0.3
The West Maui Forest Reserve is one of nine reserves on the Island of Maui that are managed by the State Department of Land and Natural Resources (DLNR)'s Division of Forestry and Wildlife. These reserves are established as multi-use land areas that incorporate various, and often competing, public uses and benefits. The management goals of the Forest Reserve System include: 1) Protect and manage forested watersheds for production of fresh water supply for public uses now and into the future; 2) Maintain biological integrity of native ecosystems; 3) Provide public recreational opportunities; and 4) Strengthen the economy by assisting in the production of high quality forest products in support of a sustainable forest industry. The primary goals of reserve are to manage the lands for sustainable game hunting, protect existing rare native biological resources and provide native and non-native timber resources.			

In addition to the individual management areas outlined above, Watershed Partnerships are another valuable component of ecosystem maintenance. Watershed Partnerships are voluntary alliances between public and private landowners who are committed to responsible management, protection, and enhancement of their forested watershed lands. There are currently nine partnerships established statewide, three of which are on Maui. Table 6-3 provides a summary of the partnership area, partners, and management goals of the West Maui Mountains Watershed Partnership.

Table 6-3. Watershed partnerships associated with the Kahoma hydrologic unit. (Source: State of Hawaii, Division of Forestry and Wildlife, 2008b)

Management Area	Year Established	Total Area (mi ²)	Area (mi ²)	Percent of Unit
West Maui Mountains Watershed Partnership	1998	78.125	3.583	62.5
The West Maui Mountains Watershed Alliance (WMMWA) is comprised of the County of Maui Department of Water Supply, Hawaii State Department of Land and Natural Resources (Division of Forestry and Wildlife, Kaanapali Land Management Corp., Kahoma Land Company L.L.C., Kamehameha Schools, Makila Land Company L.L.C., Maui Land Pineapple Company, Inc., and The Nature Conservancy of Hawaii, and Wailuku Water Company, L.L.C. The management priorities of the WMMWP include: 1) Baseline watershed forest health and threat assessments; 2) Establishment of forest health monitoring transects; 3) Fencing to control movement of feral animals, especially pigs and deer; 4) Staff control to remove pest animals from upper watershed; and 5) Pest plant control, particularly of priority weed species such as <i>Psidium cattleianum</i> . As of 2013, 19.44 miles of fencing have been built, of which 4.43 miles are pig and deer fencing (8-foot) and 14.91 miles are pig (4-foot) fencing resulting in 21,084 acres of protected lands.				

In 1974, the U.S. Fish and Wildlife Service (USFWS) initiated a National Wetlands Inventory that was considerably broader in scope than an earlier 1954 inventory that had focused solely on valuable waterfowl habitat. The inventory for Hawaii was completed in 1978 and utilized a hierarchical structure in the classification of various lands. The USFWS defines wetlands as “lands transitional between terrestrial and aquatic systems where the water table is usually at or near the surface or the land is covered by shallow water” (Cowardin et al., 1979). Nearly 11 percent of the Kahoma Hydrologic Unit is classified as seasonal, non-tidal palustrine wetlands occurring in the headwaters of the hydrologic unit (Table 6-4 and Figure 6-2). Palustrine wetlands are nontidal wetlands dominated by trees, shrubs,

persistent emergents, emergent mosses or lichens, or wetlands that occur in tidal areas where salinity due to ocean-derived salts is below 0.5 percent.

Table 6-4. Wetland classifications for Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015n)

System Type	Class	Regime	Area (mi ²)	Percent of Unit
Marine	Estuarine and Marine Deepwater	Subtidal	0.002	< 0.01
Palustrine	Freshwater Pond	Seasonal non-tidal	0.027	0.05
Palustrine	Freshwater Emergent Wetland	Seasonal non-tidal	0.022	0.04

A series of vegetation maps describing upland plant communities was prepared as part of a USFWS survey in 1976 to 1981 to determine the status of native forest birds and their associated habitats. Figure 6-3 presents the portion of the hydrologic unit surveyed for critical habitat. Nearly a half of the unit is dominated by introduced species, while almost 40 percent of the unit is dominated by native species.

The density of threatened and endangered plant species is high at elevations above 1,200 feet, resulting in the majority of the Kahoma hydrologic unit, roughly 55 percent, covered in a high or very high density of threatened and endangered plant species (Table 6-5).

Table 6-5. Density of threatened and endangered plants for Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015i)

Density	Area (mi ²)	Percent of Unit
Very High concentration of threatened and endangered species	1.651	28.5
High concentration of threatened and endangered species	1.596	27.6
Little or no threatened and endangered species	2.597	44.9

A working paper is being developed by the University of Hawaii’s Economic Research Organization (UHERO), entitled *Environmental Valuation and the Hawaiian Economy*, which discusses the use of existing measures of economic performance and alternative statistical devices to provide an economic valuation of threatened environmental resources. The paper focuses on the Koolau, Oahu watershed and illustrates three categories of positive natural capital (forest resources, shoreline resources, and water resources) against a fourth category (alien species) that degrades natural capital. In the case of the Oahu Koolau forests, a benchmark level of degradation is first defined for comparison against the current value of the Oahu Koolau system. The Oahu Koolau case study considers a hypothetical major disturbance caused by a substantial increased population of pigs with a major forest conversion from native trees to the non-indigenous *Miconia* (*Miconia calvescens*), along with the continued “creep” of urban areas into the upper watershed (Kaiser, B. et al., n.d.).

Recognizing that in the United States, the incorporation of environmental and natural resource considerations into economic measures is still very limited, the paper provides the estimated Net Present Value (NPV) for “Koolau [Oahu] Forest Amenities.” These values are presented in Table 6-7.

Following the results of the Oahu Koolau case study, some of the most valuable aspects of the forested areas are believed to be ecotourism, aesthetic pleasure, species habitat, water quality, and water quantity. Certain areas of Kahoma provide critical habitat for native forest birds, endangered plants and invertebrates in Maui.

Table 6-6. Estimated Net Present Value (NPV) for Koolau [Oahu] Forest Amenities. (Source: Kaiser, B. et al., n.d.)

Amenity	Estimated Net Present Value (NPV)	Important limitations
Ground water quantity	\$4.57 to \$8.52 billion NPV	Optimal extraction assumed.
Water quality	\$83.7 to \$394 million NPV	Using averted dredging cost estimates.
In-stream uses	\$82.4 to \$242.4 million NPV	Contingent valuation estimate for a single small fish species.
Species habitat	\$487 to \$1,434 million NPV	Contingent valuation estimate for a single small bird species.
Biodiversity	\$660,000 to \$5.5 million NPV	Average cost of listing 11 species in Koolaus.
Subsistence	\$34.7 to \$131 million NPV	Based on replacement value of pigs hunted.
Hunting	\$62.8 to \$237 million NPV	Based on fraction of hunting expenditures in state. Does not include damages from pigs to the other amenities.
Aesthetic values	\$1.04 to \$3.07 million NPV	Contingent valuation; Households value open space for aesthetic reasons.
Commercial harvests	\$600,000 to \$2.4 million NPV	Based on small sustainable extraction of koa.
Ecotourism	\$1.0 to \$2.98 billion NPV	Based on fraction of direct revenues to ecotourism activities.
Climate control	\$82.2 million	Based on replacement costs of contribution of all tropical forests to carbon sequestration.
Estimated value of joint services:	\$7.444 to \$14.032 billion	

Figure 6-2. Reserves and wetlands for the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2007b; 2015n)




Figure 6-3. Distribution of native and alien plant species, and critical habitat for plant and bird species in the Kahoma hydrologic unit. (Source: Scott et al., 1986; State of Hawaii, Office of Planning, 1996; 2004b)





Figure 6-4. Density of threatened and endangered plants in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015h)



7.0 Aesthetic Values

Aesthetics is a multi-sensory experience related to an individual's perception of beauty. Since aesthetics by definition is a subjective observation, a stream's aesthetic value cannot be determined quantitatively (Wilson Okamoto & Associates, Inc., 1983). However, there are certain elements, either within or surrounding a stream, which appeal to an observer's visual and auditory senses, such as waterfalls and cascading plunge pools. Visitors and residents can identify a point that has aesthetic value and continue to return to such a point to gain that value. Such points can potentially be identified, as mapped in Figure 5-2. However, the points listed are not exhaustive and it is beyond the scope of this report to list all potential aesthetic values.

The headwaters of Kahoma Stream originate in the West Maui Mountains, a unique ecological region home to endemic and endangered birds, plants, and insects. However, access to this region is restricted to the general public.

In a 2007 Hawaii State Parks Survey, released by the Hawaii Tourism Authority (OmniTrak Group Inc., 2007), scenic views accounted for 21 percent of park visits statewide. Other aesthetic-related motivations include viewing famous landmarks (9 percent), hiking trails and walks (7 percent), guided tour stops (6 percent), and viewing of flora and fauna (2 percent). On the island of Maui, out-of-state visitors' most common reasons to visit state parks for scenic views (26 percent) was the number one reason for visiting a park above outings with family and friends (25 percent). By contrast, residents primarily used state parks on Maui for outings with family and friends (33 percent) followed by ocean/water activity (23 percent) and then scenic views (11 percent). Overall, Maui residents were very satisfied with scenic views giving a score of 9.7 (on a scale of 1 to 10, with 10 being outstanding), with out-of-state visitors giving a score of 9.3.

8.0 Navigation

The State Water Code, Chapter 174C, HRS, includes navigation as one of nine identified instream uses; however, it fails to further define navigation. Navigational water use is largely defined as water utilized for commercial, and sometimes recreational, transportation. In the continental United States, this includes water used to lift a vessel in a lock or to maintain a navigable channel level. Under the provisions of the Clean Water Act, navigable waters also include wetlands (State of Nevada, Department of Conservation and Natural Resources, Division of Water Resources, n.d.).

Hawaii streams are generally too short and steep to support navigable uses. If recreational boating (primarily kayaks and small boats) is included under the definition of navigation, then there are only a handful of streams statewide that actually support recreational boating and even fewer that support commercial boating operations. Kauai's Wailua River is the only fresh water waterway where large boat commercial operations exist, and no streams are believed to serve as a means for the commercial transportation of goods.

The hydrologic unit of Kahoma is not known to support any instream uses of navigation.

9.0 Instream Hydropower Generation

The generation of hydropower is typically accomplished through instream dams and power generators; however, the relatively short lengths and flashy nature of Hawaii's streams often require water to be diverted to offstream power generators. In these "run-of-river" (i.e., utilizes water flow without dams or reservoirs) designs, water is diverted through a series of ditches, pipes, and penstocks to the powerplant, and then returned to the stream. Some designs call for the powerplant to be situated such that the drop of water level (head) exiting the plant can be sent to fields for crop irrigation.

There is no instream hydropower in the Kahoma hydrologic unit.

10.0 Maintenance of Water Quality

The maintenance of water quality is important due to its direct impact upon the maintenance of other instream uses such as fish and wildlife habitat, outdoor recreation, ecosystems, aesthetics, and traditional and customary Hawaiian rights. There are several factors that affect a stream's water quality, including physical, chemical, and biological attributes. The State of Hawaii Department of Health (DOH) is responsible for water quality management duties statewide. The DOH Environmental Health Administration oversees the collection, assessment, and reporting of numerous water quality parameters in three high-priority categories:

- Possible presence of water-borne human pathogens;
- Long-term physical, chemical and biological components of inland, coastal, and oceanic waters; and
- Watershed use-attainment assessments, identification of sources of contamination, allocation of those contributing sources, and implementation of pollution control actions.

The Environmental Health Administration is also responsible for regulating discharges into State waters, through permits and enforcement actions. Examples include federal National Pollutant Discharge Elimination System (NPDES) permits for storm water, and discharge of treated effluent from wastewater treatment plants into the ocean or injection wells.

Sediment and temperature are among the primary physical constituents of water quality evaluations. They are directly impacted by the amount of water in a stream. The reduction of streamflow often results in increased water temperatures, whereas higher flows can aid in quickly diluting stream contamination events. According to a book published by the Instream Flow Council, “[w]ater temperature is one of the most important environmental factors in flowing water, affecting all forms of aquatic life (Amear et al, 2004).” While this statement is true for continental rivers, fish in Hawaii are similar, but their main requirement is flowing water. Surface water temperatures may fluctuate in response to seasonal and diurnal variations, but only a few degrees Celsius in natural streams, mainly because streams in Hawaii are so short. However, temperatures in streams with concrete-lined channels, and dewatered streams, may fluctuate widely due to the vertical solar contact. Surface water temperatures may also fluctuate widely due to water column depth, channel substrate, presence of riparian vegetation, and ground water influx. Surface water also differs considerably from ground water, generally exhibiting lower concentrations of total dissolved solids, chlorides, and other major ions, along with higher concentrations of suspended solids, turbidity, microorganisms, and organic forms of nutrients (Lau and Mink, 2006). Findings of a 2004 USGS National Water Quality Assessment (NAWQA) Program report identified land use, storm-related runoff, and ground water inflow as major contributors of surface water contaminants (Anthony et al., 2004). Runoff transports large amounts of sediment from bare soil into surface water bodies, with consequences for in-stream and near-shore environments.

Water body types can be freshwater, marine, or brackish. They can be further delineated as inland fresh waters, estuaries, embayments, open coastal waters, and oceanic waters (HAR 11-54-5 to 11-54-6). Each water body type has its own numeric criteria for State of Hawaii Water Quality Standards (WQS).

Fresh waters are classified for regulatory purposes, according to the adjacent land's conservation zoning. There are two classes for the inland fresh waters. Class 1 inland waters are protected to “remain in their natural state as nearly as possible with an absolute minimum of pollution from any human-caused source.” These waters are used for a number of purposes including domestic water supply, protection of native breeding stock, and baseline references from which human-caused changes can be measured.

Class 2 inland waters are protected for uses such as recreational purposes, support of aquatic life, and agricultural water supplies.

Class 1 waters are further separated into Classes 1a and 1b. Class 1a waters are protected for the following uses: scientific and educational purposes, protection of native breeding stock, baseline references from which human-caused changes can be measured, compatible recreation, aesthetic enjoyment, and other non-degrading uses which are compatible with the protection of the ecosystems associated with waters of this class. Streams that run through natural reserves, preserves, sanctuaries, refuges, national and state parks, and state or federal fish and wildlife refuges are Class 1a. Streams adjacent to the most environmentally sensitive conservation subzone, “protective,” are Class 1b, and are protected for the same uses as Class 1a waters, with the addition of domestic water supplies, food processing, and the support and propagation of aquatic life (HAR 11-54-3). These classifications are used for regulatory purposes, restricting what is permitted on the land around receiving waters. For example, public access to Class 1b waters may be restricted to protect drinking water supplies.

Land use affects water quality because direct runoff (rainfall that flows overland into the stream) can transport sediment and its chemical contaminants into the stream. According to the U.S. Environmental Protection Agency (USEPA), “[a] TMDL or Total Maximum Daily Load is a calculation of the maximum amount of a pollutant that a waterbody can receive and still meet water quality standards, and an allocation of that amount to the pollutant's sources. Water quality standards are set by States, Territories, and Tribes. They identify the uses for each waterbody, for example, drinking water supply, contact recreation (swimming), and aquatic life support (fishing as well as ecological health), and the scientific criteria required to support those uses. A TMDL is the sum of the allowable loads of a single pollutant from all contributing point and nonpoint sources. The calculation must include a margin of safety to ensure that the waterbody can be used for the purposes the State has designated. The calculation must also account for seasonal variation in water quality. The Clean Water Act, section 303, establishes the water quality standards and TMDL programs (USEPA, 2008).”

The DOH, Environmental Health Administration maintains the State of Hawaii Water Quality Standards (WQS), a requirement under the Federal Clean Water Act (CWA) regulated by the EPA. The CWA aims to keep waters safe for plants and animals to live and people to wade, swim, and fish. Water Quality Standards are the measures that states use to ensure protection of the physical, chemical, and biological health of their waters. “A water quality standard defines the water quality goals of a water body, or portion thereof, by designating the use or uses to be made of the water and by setting criteria necessary to protect the uses (CWA §131.2).” Each state specifies its own water uses to be achieved and protected (“designated uses”), but CWA §131.10 specifically protects “existing uses”, which it defines as “...those uses actually attained in the water body on or after November 28, 1975, whether or not they are included in the water quality standards (CWA §131.3).”¹ Although the State WQS do not specify any designated uses in terms of traditional and customary Hawaiian rights, the “protection of native breeding stock,” “aesthetic enjoyment,” and “compatible recreation” are among the designated uses of Class 1 inland

¹ Existing uses as defined in the CWA should not be confused with existing uses as defined in the State Water Code, although there is some overlap and linkage between the two. Under the Water Code, if there are serious threats to or disputes over water resources, the Commission may designate a “water management area.” Water quality impairments, including threats to CWA existing uses, are factors that the Commission may consider in its designation decisions. Once such a management area is designated, people who are already diverting water at the time of designation may apply for water use permits for their “existing uses.” The Commission then must weigh if the existing use is “reasonable and beneficial.” The Water Code defines “reasonable-beneficial use” as “the use of water in such a quantity as is necessary for economic and efficient utilization, for a purpose, and in a manner which is both reasonable and consistent with the state and county land use plans and the public interest.” The relationships between a Commission existing use and a CWA existing use can help determine the appropriateness of the use and its consistency with the public interest.

waters, and “recreational purposes, the support and propagation of aquatic life, and agricultural and industrial water supplies” are among the designated uses of Class 2 inland waters. This means that uses tied to the exercise of traditional and customary Hawaiian rights that are protected by the State Constitution and the State Water Code (Section 12.0, Protection of Traditional and Customary Hawaiian Rights), including but not limited to gathering, recreation, healing, and religious practices are also protected under the CWA and the WQS as designated and/or existing uses. Therefore, the Commission’s interim IFS recommendation may impact the attainment of designated and existing uses, water quality criteria, and the DOH antidegradation policy, which together define the WQS and are part of the joint Commission and DOH obligation to assure sufficient water quality for instream and noninstream uses.

State of Hawaii WQS define: 1) the classification system for State surface waters, which assigns different protected uses to different water classes; 2) the specific numeric or narrative water quality criteria needed to protect that use; and 3) a general antidegradation policy, which maintains and protects water quality for the uses defined for a class. Quantitative and qualitative data are utilized. Numeric water quality criteria have specific concentrations (levels of pollutants) that must be attained based on water body type, e.g. fresh water stream. Qualitative standards are general narrative statements that are applicable to all State waters, such as “all waters shall be free of substances attributable to domestic, industrial, or other controllable sources of pollutants (State of Hawaii, Department of Health, 2004).” Conventional pollutants include nutrients and sediments. Toxic pollutants include pesticides and heavy metals. Indicator bacteria are utilized to assess bacterial levels. Biological assessments of aquatic communities are also included in the data collected.

Once data are gathered and evaluated for quality and deemed to be representative of the waterbody segment, a decision is made as to whether the appropriate designated uses are being attained. This set of decisions are then tabulated into a report to the EPA that integrates two CWA sections; (§) 305(b) and §303(d). This Integrated Report is federally required every even-numbered year. CWA §305(b) requires states to describe the overall water quality statewide. They must also describe the extent to which water quality provides for the protection and propagation of a balanced population of shellfish, fish, and wildlife and allows recreational activities in and on the water. Additionally, they determine whether the designated uses of a water body segment are being attained, and if not, what are the potential causes and sources of pollution. The CWA §303(d) requires states to submit a list of Water-Quality Limited Segments, which are waters that do not meet state water quality standards and those waters’ associated uses. States must also provide a priority ranking of waters listed for implementation of pollution controls, which are prioritized based on the severity of pollution and the uses of the waters. In sum, the §303(d) list leads to action.

The sources for the 2014 Integrated Report are Hawaii’s 2012 §303(d) list, plus readily-available data collected from any State water bodies over the preceding 6 years (State of Hawaii, Department of Health, 2007). Per §303(d), impaired waters are listed after review of “‘all existing and readily available water quality-related data and information’ from a broad set of data sources” (State of Hawaii, Department of Health, 2004, p.57). However, available data are not comprehensive of all the streams in the State. According to the Hawaii Administrative Rules Title 11 Chapter 54 (HAR 11-54) all State waters are subject to monitoring; however, in the most recent list published (from the 2012 list that was published in 2014), only two streams statewide had sufficient data for evaluation of whether exceedence of WQS occurred. Kahoma Stream did not appear on the 2012 List of Impaired Waters in Hawaii, Clean Water Act §303(d) but there was insufficient data for decision-making; therefore, no decision was made pertaining to the attainment of WQS or the applicable designated uses.

The 2006 Integrated Report indicates that the current WQS require the use of *Enterococci* as the indicator bacteria for evaluating public health risks in the waters of the State; however, no new data were available for this parameter in inland waters. As mentioned in Section 5.0, Outdoor Recreational Activities, DOH


maintains WQS for inland recreational waters based on the geo-mean statistic of *Enterococci*: 33 colony-forming units per 100 mL of water or a single-sample maximum of 89 colonies per 100 mL. This is for full-body contact (swimming, jumping off cliffs into waterfall pools, etc.). If *Enterococci* count exceeds those values, the water body is considered to be impaired. DOH Clean Water Branch efforts have been focused on coastal areas (State of Hawaii, Department of Health, 2006, Chapter II, p.20). The marine recreational zone, which extends from the shoreline seaward to 1,000 feet from shore, requires an *Enterococci* geo-mean of less than 7 colony-forming units per 100 mL of water to protect human health (HAR 11-54-8.)

The 2012 Integrated Report also states: “Public health concerns may be underreported. *Leptospirosis* is not included as a specific water quality standard parameter. However, all fresh waters within the state are considered potential sources of *Leptospirosis* infection by the epidemiology section of the Hawaii State Department of Health. No direct tests have been approved or utilized to ascertain the extent of the public health threat through water sampling. Epidemiologic evidence has linked several illness outbreaks to contact with fresh water, leading authorities to issue blanket advisories for all fresh waters of the state (State of Hawaii, Department of Health, 2007, Chapter II, p.3).”

Sections of Kahoma Stream are classified as Class 1a inland waters from its headwaters to the boundary of the forest reserve as the surrounding land is in the conservation subzone “protective.” Below this, the stream is classified as Class 1b inland waters to the coastline. It should be noted that the conservation subzone map utilized for this interpretation is general and elevations are not exact. It should also be noted that there is no direct relationship between elevation and attainment of water quality standards.

Marine water body types are delineated by depth and coastal topography. Open coastal waters are classified for protection purposes from the shoreline at mean sea level laterally to where the depth reaches 100 fathoms (600 feet). Marine water classifications are based on marine conservation areas. The objective of Class AA waters is that they “remain in their natural pristine state as nearly as possible with an absolute minimum of pollution or alteration of water quality from any human-caused source or actions.” Class A waters are protected for recreational purposes and aesthetic enjoyment; and protection of fish, shellfish, and wildlife. Discharge into these waters is permitted under regulation. The marine waters at the mouth of the Kahoma hydrologic unit are Class A waters. Figure 10-1 shows the Kahoma hydrologic unit, including inland and marine (coastal) water classifications.

Figure 10-1. Water quality standards for the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015e; 2008). The classifications are general in nature and should be used in conjunction with Hawaii Administrative Rules, Chapter 11-54, Water Quality Standards.



11.0 Conveyance of Irrigation and Domestic Water Supplies

Under the State Water Code, the conveyance of irrigation and domestic water supplies to downstream points of diversion is included as one of nine listed instream uses. The thought of a stream as a conveyance mechanism for noninstream purposes almost seems contrary to the concept of instream flow standards. However, the inclusion of this instream use is intended to ensure the availability of water to all those who may have a legally protected right to the water flowing in a stream. Of particular importance in this section is the diversion of surface water for domestic purposes. In its August 2000 decision on the Waiahole Ditch Combined Contested Case Hearing, the Hawaii Supreme Court identified domestic water use of the general public, particularly drinking water, as one of, ultimately, four trust purposes.

Neither the State nor the County keeps a comprehensive database of households whose domestic water supply is not part of a municipal system (i.e. who use stream and / or catchment water). The County of Kauai Department of Water Supply does not have data for water users who are not on the county system and may be using catchment or surface water for domestic use. The State of Hawaii Department of Health Safe Drinking Water Branch administers Federal and State safe drinking water regulations to public water systems in the State of Hawaii to assure that the water served by these systems meets State and Federal standards. Any system which services 25 or more people for a minimum of 60 days per year or has at least 15 service connections is subject to these standards and regulations. Once a system is regulated by the Safe Drinking Water Branch, the water must undergo an approved filtration and disinfection process when it has been removed from the stream. It would also be subject to regulatory monitoring. However, there are no private water systems in the Kahoma hydrologic unit regulated by the DOH, Safe Drinking Water branch. Maui County operates a municipal water treatment facility that uses surface water from the Kanaha Stream, a tributary of Kahoma Stream. This facility (Lahaina Surface Water Treatment Facility) processes an average of 1.76 mgd, with the capacity to process up to 3 mgd.

12.0 Protection of Traditional and Customary Hawaiian Rights

The maintenance of instream flows is important to the protection of traditional and customary Hawaiian rights, as they relate to the maintenance of stream resources (e.g., hihiwai, opae, oopu) for gathering, recreation, and the cultivation of taro. Article XII, Section 7 of the State Constitution addresses traditional and customary rights: “The State reaffirms and shall protect all rights, customarily and traditionally exercised for subsistence, cultural and religious purposes and possessed by ahupua‘a tenants who are descendants of native Hawaiians who inhabited the Hawaiian Islands prior to 1778, subject to the right of the State to regulate such rights.” Case notes listed in this section indicate, “Native Hawaiian rights protected by this section may extend beyond the ahupua‘a in which a native Hawaiian resides where such rights have been customarily and traditionally exercised in this manner. 73 H.578, 837 P.2d 1247.”

It is difficult to fully represent in words the depth of the cultural aspects of streamflow, including traditions handed down through the generations regarding gathering, ceremonial and religious rites, and the ties to water that are pronounced in Hawaiian legend and lore. “There is a great traditional significance of water in Hawaiian beliefs and cultural practices...The flow of water from mountain to sea is integral to the health of the land. A healthy land makes for healthy people, and healthy people have the ability to sustain themselves (Kumu Pono Associates, 2001b, p.II:8).”

Taro cultivation is addressed in this section of the report as well as section 14. This is because instream flow standards take into account both social and scientific information. For sociological and cultural purposes, taro cultivation can be considered an instream use as part of the “protection of traditional and customary Hawaiian rights,” that is specifically listed as an instream use in the Water Code. Taro cultivation can also be considered a noninstream use since it removes water from a stream (even if water from taro loi is later returned to the stream). It could be argued that for scientific analysis, taro cultivation is an instream use since taro loi provide habitat for stream biota, but because the water is physically taken out of the stream, it is also a noninstream use. Another way to look at the approach of indentifying taro cultivation as both instream and noninstream uses is that when the Commission addresses taro cultivation as an instream use, it is generally in the context of traditional and customary Hawaiian rights; whereas when the Commission addresses taro cultivation as a noninstream use, it is approaching the issue from the aspects of agriculture and water use.

In ancient Hawaii, the islands (*moku*) were subdivided into political subdivisions, or ahupuaa, for the purposes of taxation. The term ahupuaa in fact comes from the altar (*ahu*) that marked the seaward boundary of each subdivision upon which a wooden head of a pig (*puaa*) was placed at the time of the *Makahiki* festival when harvest offerings were collected for the rain god and his earthly representative (Handy et al., 1972). Each ahupuaa had fixed boundaries that were usually delineated by natural features of the land, such as mountain ridges, and typically ran like a wedge from the mountains to the ocean thus providing its inhabitants with access to all the natural resources necessary for sustenance. The beach, with its fishing rights, were referred to as *ipu kai* (meat bowl), while the upland areas for cultivation were called *umeke ai* (poi container hung in a net) (Handy et al., 1972). As noted earlier in Section 6.0, Maintenance of Ecosystems, Western concepts of ecosystem maintenance and watersheds are similar to the Hawaiian concept of ahupuaa, and so the Commission’s surface water hydrologic units often coincide with or overlap ahupuaa boundaries. The hydrologic unit of Kahoma includes many ahupuaa as shown in Figure 12-2. The ahupuaa boundaries are delineated based on the USGS Digital Line Graphs. These boundaries may be different from the information listed on legal documents such as deeds.

An appurtenant water right is a legally recognized right to a specific amount of surface freshwater – usually from a stream – on the specific property that has that right. This right traces back to the use of water on a given parcel of land at the time of its original conversion into fee simple lands: When the land allotted during the 1848 Mahele was confirmed to the awardee by the Land Commission and/or when the Royal Patent was issued based on such award, the conveyance of the parcel of land carried with it the appurtenant right to water if water was being used on that land at or shortly before the time of the Mahele (State of Hawaii, Commission on Water Resource Management, 2007).

An appurtenant right is different from a riparian right, but they are not mutually exclusive. Riparian rights are held by owners of land adjacent to a stream. They and other riparian landowners have the right to reasonable use of the stream's waters on those lands. Unlike riparian lands, the lands to which appurtenant rights attach are not necessarily adjacent to the freshwater source (i.e., the water may be carried to the lands via auwai or ditches), but some pieces of land could have both appurtenant and riparian rights.

Appurtenant rights are provided for under the State Water Code, HRS §174C-101, Section (c) and (d) as follows:

- Section (c). Traditional and customary rights of ahupuaa tenants who are descendants of native Hawaiians who inhabited the Hawaiian Islands prior to 1778 shall not be abridged or denied by this chapter. Such traditional and customary rights shall include, but not be limited to, the cultivation or propagation of taro on one's own kuleana and the gathering of hihiwai, opae, oopu, limu, thatch, ti leaf, aho cord, and medicinal plants for subsistence, cultural, and religious purposes.
- Section (d). The appurtenant water rights of kuleana and taro lands, along with those traditional and customary rights assured by this section, shall not be diminished or extinguished by a failure to apply for or to receive a permit under this chapter.

The exercise of an appurtenant water right is still subject to the water use permit requirements of the Water Code, but there is no deadline to exercise that right without losing it, as is the case for correlative and riparian rights, which must have been exercised before designation of a water management area.

In August 2000, the Hawaii Supreme Court issued its decision in the Waiahole Ditch Combined Contested Case Hearing, upholding the exercise of Native Hawaiian and traditional and customary rights as a public trust purpose. These rights are described in the Commission's 2007 *Water Resource Protection Plan – Public Review Draft*, incorporating a later revision¹ as follows:

Appurtenant water rights are rights to the use of water utilized by parcels of land at the time of their original conversion into fee simple lands i.e., when land allotted by the 1848 Mahele was confirmed to the awardee by the Land Commission and/or when the Royal Patent was issued based on such award, the conveyance of the parcel of land carried with it the appurtenant right to water.² The amount of water under an appurtenant right is the amount that was being used at the time of the Land Commission award and is established by cultivation methods that approximate the methods utilized at the time of the Mahele, for example, growing wetland taro.³ Once established, future uses are not limited to the cultivation of traditional products approximating

¹ Although the final Water Resource Protection Plan had not been printed as of the date of this report, most edits had already been incorporated into the latest version, which the Commission utilized for this report.

² 54 Haw. 174, at 188; 504 P.2d 1330, at 1339.

³ 65 Haw. 531, at 554; 656 P.2d 57, at 72.

those utilized at the time of the Mahele⁴, as long as those uses are reasonable, and if in a water management area, meets the State Water Code's test of reasonable and beneficial use ("the use of water in such a quantity as is necessary for economic and efficient utilization, for a purpose, and in a manner which is both reasonable and consistent with the State and county land use plans and the public interest"). As mentioned earlier, appurtenant rights are preserved under the State Water Code, so even in designated water management areas, an unexercised appurtenant right is not extinguished and must be issued a water use permit when applied for, as long as the water use permit requirements are met [Figure 12-1].

The Hawaii Legislative Session of 2002 clarified that the Commission is empowered to "determine appurtenant rights, including quantification of the amount of water entitled to by that right," (HRS §174C-5(15)). In those cases where a Commission decision may affect an appurtenant right, it is the claimant's duty to assert the appurtenant right and to gather the information required by the Commission to rule on the claim. The Commission is currently in the process of developing a procedural manual to aid in the understanding and assembling of information to substantiate an appurtenant rights claim.

In accordance with the State Water Code and the Supreme Court's decision in the Waiahole Ditch Combined Contested Case Hearing, the Commission is focused on the assertion and exercise of appurtenant rights as they largely relate to the cultivation of taro. Wetland kalo or taro (*Colocasia esculenta* (L.) Schott) is an integral part of Hawaiian culture and agricultural tradition. The preferred method of wetland taro cultivation, where terrain and access to water permitted, was the construction of loi (flooded terraces) and loi complexes. These terraces traditionally received stream water via carefully engineered open channels called auwai. The auwai carried water, sometimes great distances, from the stream to the loi via gravity flow. In a system of multiple loi, water may either be fed to individual loi through separate little ditches if possible, or in the case of steeper slopes, water would overflow and drain from one loi to the next. Outflow from the loi may eventually be returned to the stream.

The loi also served other needs including the farming of subsidiary crops such as banana, sugar cane, and ti plants that were planted on its banks, and the raising of fish such as oopu, awa, and aholehole within the waters of the loi itself. At least 85 varieties of taro were collected in 1931, each of which varied in color, locale, and growing conditions. The water needs of taro under wet conditions depend upon: 1) climate; 2) location and season (weather); 3) evaporation rate; 4) soil type; 5) ground water hydrology; 5) water temperature; and 6) agronomic conditions (crop stage; planting density and arrangement; taro variety; soil amendment and fertilization regime; loi drainage scheme; irrigation system management; and weed, pest, and disease prevalence and management).

The Commission conducted a cursory assessment of tax map key parcels to identify their associated Land Commission Awards, in an attempt to identify the potential for future appurtenant rights claims within the hydrologic unit of Kahoma. Table 12-1 presents the results of the Commission's assessment. Of particular importance is the presence of terracing and oral testimony that indicates loi kalo existed as recently as the 1980s in the upper reaches of Kahana, immediately below the current stream diversion (Figure 12-3). The series of TMK parcels and associated land commission awards suggest extensive cultivation of the riparian areas of both Kanaha and Kahoma streams.

⁴ *Peck v Bailey*, 8 Haw. 658, at 665 (1867).

Figure 12-1. Generalized process for determining appurtenant water rights. This process is generalized and may not fully explain all possible situations. It does not apply to Hawaiian Homes Lands. If you are Native Hawaiian you may have other water rights.




Figure 12-2. Land Commission Awards and associated tax map key parcel numbers for the Kahoma hydrologic unit, Maui. [LCA is Land Commission Award; Gr. is Grant; por. is portion; and G.L. is Government Lease]

LCA/Gr	TMK	Landowner	Notes
LCA 11216:15	246018007	State DLNR	Claimant: Kekauonohi, Mataio Kahoma Stream Ag/Habitation Complex: 8 terraces & 1 enclosure on north side of stream
LCA 3422 B:1; Gr 1678	246018007	State DLNR	Claimant: Kekahuna
LCA 7713:25	246017001	Keahi, William	Claimant: Kamamalu, Victoria
Gr 723	246017001	Keahi, William	Claimant: Ua, LS
LCA 6711:1; Gr 8218	246017001	Keahi, William	Claimant: Puuki
LCA 6391:2	246017014	Michel, Hans	Claimant: Kekualele
LCA 10968:2	246017002	Ralar, Nohealani Holuamoku	Claimant: Wahie
LCA 486 C:3	246017004	Michel, Hans	Claimant: Uluoā
LCA 2777	246017004	Michel, Hans	Claimant: Kalola
LCA 2320:3	246017004	Michel, Hans	Claimant: Mere
LCA 58:2.2	246017888		Claimant: Mapu
LCA 7587:2.3	246017006	Michel, Hans	Claimant: Keaweiwi
LCA 4320:2	246017006	Michel, Hans	Claimant: Kaua
LCA 3424 B:3	246017008	Michel, Hans	Claimant: Kaleleiki
LCA 11216:15.3	246017009	Michel, Hans	Claimant: Kekauonohi
LCA 962:3	246017010	Michel, Hans	Claimant: Armas, Joaquin
LCA 6787:2	246017010	Michel, Hans	Claimant: Hanaumua
LCA 11216:15.2	246017011	Adams, Renee K	Claimant: Kekauonohi
LCA 3702:2.1	246017012	Michel, Hans	Claimant: Malo, Davida
LCA 10667:3	246017012	Michel, Hans	Claimant: Pīkanele
LCA 8452:4.3	246017012	Michel, Hans	Claimant: Keohokalole, A
LCA 5017:2	246016003	Michel, Hans	Claimant: Kauhi
LCA 11086:5	246018005	State DLNR	Claimant: Kaiheekai, J H
LCA: 3425 B:2	245016007	Michel, Hans	Claimant: Alu
LCA 502:3	245016007	Michel, Hans	Claimant: Pupuka
LCA 10474:8.4	245016003	Michel, Hans	Claimant: Namauu
LCA 7713:27	245016003	Michel, Hans	Claimant: Kamamalu, Victoria
LCA 9795 G:3	245016003	Michel, Hans	Claimant: Ua
LCA 8520:3	246018006	State DLNR	Claimant: Kaeo, Iosua
LCA 355:3	245016003	Michel, Hans	Claimant: Keopenui
LCA 7632	245016001	Michel, Hans	Claimant: Kahookano, I
LCA 6412:2	245016009	Kahoma Land LLC	Claimant: Kala
LCA 8559 B:24.2	245016003	Michel, Hans	Claimant: Lunalilo, William C
LCA 525:2	245016003	Michel, Hans	Claimant: Kahiamoe
LCA 5593:3	245016003	Michel, Hans	Claimant: Kapule
LCA 355:4	245016003	Michel, Hans	Claimant: Keopenui
LCA 6498:1	245016010	Kahoma Land LLC	Claimant: Kokio
LCA 8520:2.2	245016009	Kahoma Land LLC	Claimant: Kaeo, Iosua

Table 12-1. [continued]

LCA/Gr	TMK	Landowner	Notes
LCA 6412:1	245016009	Kahoma Land LLC	Claimant: Kala
LCA 7724:2	245015008	Kahoma Land LLC	Claimant: Poholapu
LCA 6541	245016008	Pedro, Sarah H	Claimant: Haia
LCA 8559 B:26.1	245016888	n/a	Claimant: Lunalilo, William C
LCA 6413	245017007	Kahoma Land LLC	Claimant: Kaaukai
LCA 6620:1	245016009	Kahoma Land LLC	Claimant: Napahi
LCA 8559 B:26.2	245016888	n/a	Claimant: Lunalilo, William C
LCA 3421 B:3	245017005	Kahoma Land LLC	Claimant: Kaaa
LCA 7724:1	245017005	Kahoma Land LLC	Claimant: Poholapu
LCA 8559 B:26.3	245017002	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 7591	245017002	Kahoma Land LLC	Claimant: Kalakoa
LCA 8021:1	245017002	Kahoma Land LLC	Claimant: Aha
LCA 6621	245017002	Kahoma Land LLC	Claimant: Nalehu
LCA 10579	245016888	n/a	Claimant: Opunui
LCA 8559 B:26.4	245017005	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 8021:2	245017005	Kahoma Land LLC	Claimant: Aha
LCA 11149:2	245017005	Kahoma Land LLC	Claimant: Kuhua
LCA 8559 B:26.5	245017002	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 8021:3	245017005	Kahoma Land LLC	Claimant: Aha
LCA 8559 B:26.6	245017002	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 8515:3.2	245021001	BP Bishop Trust Estate	Claimant: Lahilahi, Gini
LCA 8559 B:26.7	245018011	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 9812 D	245018011	Kahoma Land LLC	Claimant: Kanalu
LCA 11216:19.2	245021022	State DLNR	Claimant: Kekauonohi, M
LCA 466:1	245018011	Kahoma Land LLC	Claimant: Keone
LCA 447 F:1	245018011	Kahoma Land LLC	Claimant: Keliipio
LCA 8559 B:26.8	245018011	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 466:2	245018011	Kahoma Land LLC	Claimant: Keone
LCA 8559 B:26.9	245018009	Dean, Joshua I	Claimant: Lunalilo, William C
LCA 6061:5	245018011	Kahoma Land LLC	Claimant: Hanemo
LCA 6052:3	245018011	Kahoma Land LLC	Claimant: Waihoikaea
LCA 7582	245021002	Kahoma Land LLC	Claimant: Kipa, Eseta
LCA 8559 B:26.10	245021002	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 486:3	245018013	Kahoma Land LLC	Claimant: Ku, W
LCA 6061:6	245018008	Kahoma Land LLC	Claimant: Hanemo
LCA 10474:8.2	245018008	Kahoma Land LLC	Claimant: Namauu
LCA 470:3	245018008	Kahoma Land LLC	Claimant: Kuokoa
LCA 9795:3	245018008	Kahoma Land LLC	Claimant: Kaaumaiewa
LCA 9793:4	245018008	Kahoma Land LLC	Claimant: Mumuku
LCA 9795:4	245018008	Kahoma Land LLC	Claimant: Kaaumaiewa
LCA 470:4	245018007	Power, Mary	Claimant: Kuokoa

Table 12-1. [continued] Land Commission Awards and associated tax map key parcel numbers for the Kahoma hydrologic unit, Maui. [LCA is Land Commission Award; Gr. is Grant; por. is portion; and G.L. is Government Lease; L.C.App. is Land Court Application]

LCA/Gr	TMK	Landowner	Notes
LCA 5912:4	245018008	Kahoma Land LLC	Claimant: Pahia, Simeona
LCA 5483:5	245018001	Kahoma Land LLC	Claimant: Kaeo, I
LCA 9950:1	245018001	Kahoma Land LLC	Claimant: Leleku
LCA 10431:1	245018001	Kahoma Land LLC	Claimant: Nipoa
LCA 7713:26	245021001	BP Bishop Trust Estate	Claimant: Kamamalu, Victoria
LCA 9780 B:1	245018004	BP Bishop Trust Estate	Claimant: Kaumiumi
LCA 7713:25	245018006	Kahoma Land LLC	Claimant: Kamamalu, Victoria
Gr 962:1	245018003	BP Bishop Trust Estate	Claimant: Puhalahua
LCA 9950:2	245018001	Kahoma Land LLC	Claimant: Leleku
LCA 474:2	245018001	Kahoma Land LLC	Claimant: Kalei
LCA 7716:11.2	245019017	BP Bishop Trust Estate	Claimant: Keelikolani, Ruth
LCA 7582:3	245019015	Kahoma Land LLC	Claimant: Kipa, Eseta
LCA 10431:2	245019015	Kahoma Land LLC	Claimant: Nipoa
LCA 1001:2	245019014	Kahoma Land LLC	Claimant: Keaweluaole
LCA 2538:4	245019014	Kahoma Land LLC	Claimant: Umiumi
LCA 2538:3	245019014	Kahoma Land LLC	Claimant: Umiumi
LCA 6484:1	245019010	Kahoma Land LLC	Claimant: Kanae
LCA 2924:3	245019006	Kahoma Land LLC	Claimant: Kailaa, Kale
LCA 488	245019018	Kahoma Land LLC	Claimant: Kaai, B
LCA 1001:3	245019006	Kahoma Land LLC	Claimant: Keaweluaole
LCA 2762:1	245019006	Kahoma Land LLC	Claimant: Nawaakoa
LCA 9780 B:2	245019006	Kahoma Land LLC	Claimant: Kaumiumi
LCA 5107:1	245019006	Kahoma Land LLC	Claimant: Keaka
LCA 6403:1	245019888	n/a	Claimant: Puu
LCA 5107:2	245019002	Kahoma Land LLC	Claimant: Keaka
LCA 2871:1	245020006	Kahoma Land LLC	Claimant: Kaneoalii
LCA 4804:3	245020005	Kahoma Land LLC	Claimant: Nui
LCA 2871:2	245020003	Kahoma Land LLC	Claimant: Kaneoalii
LCA 6484:2	245020002	Kahoma Land LLC	Claimant: Kanae
LCA 277:2.2	245015010	Kahoma Land LLC	Claimant: Lunalilo, William C
LCA 5006	245015888	n/a	Claimant: Kalena
LCA 6432:2	245015010	Kahoma Land LLC	Claimant: Kaninau
LCA 6408	245015010	Kahoma Land LLC	Claimant: Kaleikini, J
LCA 393:2	245015888	n/a	Claimant: Kekuelike
LCA 8452:4.4	245010008	Pioneer Mill Co	Claimant: Keohokalole
LCA 7713:25	245010008	Pioneer Mill Co	Claimant: Kamamalu, Victoria
LCA 9795 B:1	245010008	Pioneer Mill Co	Claimant: Kaaua
LCA 4801:2	245010008	Pioneer Mill Co	Claimant: Laanui, G
LCA 6061:4	245021003	State HHFDC	Claimant: Hanemo
LCA 11227:5	245010060	Mate Maui LLC	Claimant: Nalimu
Gr 1891:8	245010063	unknown	Claimant: Baldwin, Dwight

Table 12-1. [continued]

LCA/Gr	TMK	Landowner	Notes
LCA 6760 B:6	245010063	unknown	Claimant: Hanalei
LCA 6849:1	245010008	Pioneer Mill Co	Claimant: Nahale
Gr 1891:9	245010008	Pioneer Mill Co	Claimant: Baldwin, Dwight
Gr 2998	245010005	West Maui Land Co	Claimant: Jones, William Ap
LCA 312:1	245010008	Pioneer Mill Co	Claimant: Keaweiki, Timoteo
LCA 3702 & 6659:2	245010008	Pioneer Mill Co	Claimant: Malo, Davida
LCA 6073:1	245010025	Emerald Plaza Center	Claimant: Halepaka & Aui
Gr 1891:7	245010005	West Maui Land Co	Claimant: Baldwin, Dwight
LCA 4760:1	245010005	West Maui Land Co	Claimant: Lelehu
LCA 6424:2	245010005	West Maui Land Co	Claimant: Kanehoewaa
LCA 7582:1	245024051	Richardson, Eric	Claimant: Kipa, Eseta
LCA 7762	245024067	Catugal, Stanford	Claimant: Kaninaualii
Gr 1875	245009007	Pioneer Mill Co	Claimant: Shaw, George
LCA 11149:1	245009007	Pioneer Mill Co	Claimant: Kuhua
LCA 11216:19.1	245007029	Kaanapali Development Corp	Claimant: Kekauonohi, M
LCA 281 B:1	245007026	Conroy, Harold W	Claimant: Ali
LCA 5116:2	245007024	Vulcan Hawaii Inc	Claimant: Hanae, Samuela
LCA 5593:1	245007005	State HCDCH	Claimant: Kapule
LCA 355:1	245007999	Road	Claimant: Keopenui
LCA 7715:3	245007999	Road	Claimant: Kamehamea, Lot
LCA 486:1	245039999	Road	Claimant: Ku, W
10474:8.1	245003013	3900 Corp	Claimant: Namauu
LCA 6061:2	245003013	3900 Corp	Claimant: Hanemo
LCA 3425 B:3	245039999	Road	Claimant: Alu
LCA 9795 G:2	245008001	The Weinberg Foundation	Claimant: Ua
LCA 520:2	245008001	The Weinberg Foundation	Claimant: Ii, Daniela
LCA 3421 B:1	245008001	The Weinberg Foundation	Claimant: Kaaa
LCA 3702:2.3	245011001	Opukea At Lahaina Condo Master	Claimant: Malo, Davida
LCA 8515:3.1	245008001	The Weinberg Foundation	Claimant: Lahilahi, Gini
LCA 6061:1	245008001	The Weinberg Foundation	Claimant: Hanemo
LCA 6052:1	245008001	The Weinberg Foundation	Claimant: Waihokaea & Kenui
LCA 277:4	245004010	Mala Partners LC	Claimant: Lunalilo, William C
LCA 6073:2	245004022	Ellen Trust	Claimant: Halepaka & Ani
LCA 6052:2	245004002	Puunoa Beach Estates	Claimant: Waihokaea & Kenui
LCA 327:2.1	245004002	Puunoa Beach Estates	Claimant: Kaauwai, Z
LCA 11224	245004002	Puunoa Beach Estates	Claimant: Kalalau
Gr 962:2	245004051	Aoao of Puunoa Beach Estates	Claimant: Puhalahua
LCA 333	245003037	Gay Family Trust	Claimant: Kaahanui
LCA 10474:8.1	245003006	Mcdonald, Mark K	Claimant: Namauu
LCA 486:2	245003020	Gates, Kevin	Claimant: Ku, W
LCA 6760 B:5	245010008	Pioneer Mill Co	Claimant: Hanalei

Table 12-1. [continued]

LCA/Gr	TMK	Landowner	Notes
LCA 6849:2	245011007	County of Maui	Claimant: Nahale
LCA 6760 B:4	245011007	County of Maui	Claimant: Hanalei
LCA 7748	245011002	unknown	Claimant: Keahilaweiki
LCA 7750:1	245011002	unknown	Claimant: Kuikawai
Gr 1891:2	245011002	unknown	Claimant: Baldwin, Dwight
LCA 5904:1	245011003	Property Development Centers	Claimant: Polea
LCA 6760 B:1	245005009	Cannery Commercial	Claimant: Hanalei
LCA 6760 B:2	245011004	Lahaina Cannery Shops	Claimant: Hanalei
LCA 286:6	245011002	unknown	Claimant: Keawa
LCA 3702 & 6659:1	245011002	unknown	ClaimantL Malo, Davida
LCA 286:5	245011002	unknown	Claimant: Keawa
LCA 2320:2	245011002	unknown	Claimant: Mere
Gr 1891:3	245011002	unknown	Claimant: Baldwin, Dwight
LCA 10532:2	245011008	Lahaina Gateway Commercial	Claimant: Naolulo
LCA 2320:1	245012999	road	Claimant: Mere
LCA 486 C:2	245012999	road	Claimant: Uluoa
LCA 6850	245005011	Cannery Commerical	Claimant: Mahuka
Gr 1891:1	245005011	Cannery Commerical	Claimant: Baldwin, Dwight
LCA 11216:16	245012999	road	Claimant: Kekauonohi, Mataio
LCA 7749	245012999	road	Claimant: Kanananui
LCA 303-307:4	245012048	Truong, Henry	Claimant: Kaukamauna, E
LCA 5483:1	245012031	White, Alohalinda	Claimant: Kaeo, I
LCA 303-307:5	245012037	Taufa, Makamoeafi	Claimant: Kaukamauna, E
Gr 7855	245012001	Macdonald, Lucille	Claimant: Lahaina Ag Co
Gr 1871	245012056	Azbill, Gerald	Claimant: Kekai
LCA 477 F:2	245013027	Lahaina Roads	Claimant: Keliipio
LCA 5483:2	245012999	road	Claimant: Kaeo, I

Taro Production

In 2002, the State Office of Hawaiian Affairs cosponsored a “No Ka Lo‘i Conference”, in the hopes of bringing together taro farmers from around the state to share knowledge on the cultivation of taro. An outcome of the conference was an acknowledgement that farmers needed to better understand the water requirements of their taro crops to ensure and protect their water resource interests. The result of this effort was a 2007 USGS wetland kalo water use study, prepared in cooperation with the State Office of Hawaiian Affairs, which specifically examined flow and water temperature data in a total of 10 cultivation areas on four islands in Hawaii.

The study reiterated the importance of water temperature in preventing root rot. Typically, the water in the taro loi is warmer than water in the stream because of solar heating. Consequently, a taro loi needs continuous flow of water to maintain the water temperature at an optimum level. Multiple studies cited in Gingerich, et al., 2007, suggest that water temperature should not exceed 77°F (25°C). Low water temperatures slow taro growth, while high temperatures may result in root rot (Penn, 1997). When the

flow of water in the stream is low, possibly as a result of diversions or losing reaches, the warmer water in the taro loi is not replaced with the cooler water from the stream at a quick enough rate to maintain a constant water temperature. As a result, the temperature of the water in the taro loi rises, triggering root rot.

The 2007 USGS study noted that “although irrigation flows for kalo cultivation have been measured with varying degrees of scientific accuracy, there is disagreement regarding the amount of water used and needed for successful kalo cultivation, with water temperature recognized as a critical factor. Most studies have focused on the amount of water consumed rather than the amount needed to flow through the irrigation system for successful kalo cultivation (Gingerich, et al., 2007).” As a result, the study was designed to measure the throughflow of water in commercially viable loi complexes, rather than measuring the consumption of water during taro growth.

Because water requirements for taro vary with the stage of maturity of the plants, all the cultivation areas selected for the study were at approximately the same stage (i.e. near harvesting, when continuous flooding is required). Temperature measurements were made every 15 minutes for approximately 2 months. Flow measurements were collected at the beginning and the end of that period. Data were collected during the dry season (June – October), when water requirements for cooling kalo are higher. Surface water temperatures generally begin to rise in April and remain elevated through September, due to increased solar heating. Water inflow temperature was measured in 17 loi complexes, and only three had inflow temperatures rising above 27°C (the threshold temperature above which wetland kalo is more susceptible to fungi and associated rotting diseases).

The average and median inflows from all 10 cultivation areas studied are listed in Table 12-4 below. The study indicated that the “values are consistent with previously reported inflow and are significantly higher than values generally estimated for consumption during kalo cultivation.” It should also be noted that farmers were interviewed during field visits; most “believed that their supply of irrigation water was insufficient for proper kalo cultivation.”

The study results are presented in Table 12-5 (discharge measurements) and Table 12-6 (water-temperature statistics).

Table 12-1. Summary of water use calculated from loi and loi complexes by island, and the entire state. (Source: Gingerich et al., 2007, Table 10)

[gad = gallons per acre per day; na = not available]

Island	Complex			Loi				
	Number	Average water use (gad)	Average windward water use (gad)	Average leeward water use (gad)	Number	Average water use (gad)	Average windward water use (gad)	Average leeward water use (gad)
Kauai	6	120,000	97,000	260,000	2	220,000	220,000	na
Oahu	5	310,000	380,000	44,000	4	400,000	460,000	210,000
Maui	6	230,000	230,000	na	na	na	na	na
Hawaii	2	710,000	710,000	na	na	na	na	na
Average of all measurements		260,000	270,000	150,000		350,000	370,000	210,000
Median of all measurements		150,000	150,000	150,000		270,000	320,000	210,000

Table 12-2. Water-temperature statistics based on measurements collected at 15-minute intervals for loi complexes on the island of Maui. (Source: Gingerich et al., 2007, Table 7)
[°C = degrees Celsius; na = not applicable]

Geographic designation	Area	Station	Period of record	Temperature (°C)			Temperature measurements greater than 27°C (percent)
				Mean	Range	Mean daily range	
Windward	Waihee	Ma08A-CI	7/29/2006 - 9/22/2006	21.6	19.9 - 24.0	2.0	0.0
		Ma08B-CIL	7/29/2006 - 9/22/2006	24.9	20.3 - 34.0	7.6	25.4
		Ma08B-CO	7/29/2006 - 9/22/2006	25.5	20.0 - 35.5	5.7	27.0
Windward	Wailua (Lakini)	Ma09-CIT	7/30/2006 - 9/21/2006	20.7	18.5 - 23.4	2.3	0.0
		Ma09-CO	7/30/2006 - 9/21/2006	23.2	18.4 - 31.7	7.4	16.9
Windward	Wailua	Ma10-CI	7/30/2006 - 9/21/2006	22.5	20.5 - 25.9	1.9	0.0
Windward	Wailua (Waikani)	Ma11-CI	7/30/2006 - 9/21/2006	22.2	21.0 - 24.0	0.7	0.0
		Ma11-CO	7/30/2006 - 9/21/2006	26.1	22.1 - 31.8	3.3	29.1
Windward	Keanae	Ma12-CI	7/31/2006 - 9/21/2006	20.0	19.0 - 21.9	1.0	0.0
		Ma12-CO	equipment malfunction	na	na	na	na

Historical uses can also provide some insight into the protection of traditional and customary Hawaiian rights. Handy et al., in *Native Planters of Old Hawaii* (1972), provide a limited regional description as follows:

Lahaina District was a favorable place for the high chiefs of Maui and their entourage for a number of reasons: the abundance of food from both land and sea; its equable climate and its attractiveness as a place of residence; it had probably the largest concentration of population, with its adjoining areas of habitation; easy communication with the other heavily populated areas...; and its propinquity to Lanai and Molokai. [p. 492]

Lahaina's main taro lands, on the lower slopes running up to the west side of Puu Kukui, were watered by two large streams, Kanaha and Kahoma, which run far back into deep valleys whose sides were too precipitous for terracing.

As quoted in Handy et al. (1972; p. 493), Menzies wrote in 1973:

We observed rugged banks of a large rivulet that...watered with great neatness and industry. Even the shelving cliffs of rock were planted with esculent roots, banked in and watered by the most ingenious engineer.

...we found scattered along shore on a low tract of land that was neatly divided into little fields and laid out in the highest state of cultivation and improvement by being planted in the most regular manner with the different esculent roots and useful vegetables of the country, and watered at pleasure by aqueducts that ran here and there along the banks intersecting the fields.

Also quoted in Handy et al. (1972; p. 493), Argo wrote in 1823:

The environs of Lahaina are like a garden. It would be difficult to find a soil more fertile, or a people who can turn it to greater advantage; little pathways sufficiently raised, and kept in excellent condition...frequently divided by trenches, through which a fresh and limpid stream flows tranquilly, giving life to the plantations.

The space cultivated by the natives of Lahaina is about three leagues in length, and one in its greatest breadth. Beyond this all is dry and barren.

Without delving into the extensive archive of literature (refer to Kumu Pono Associates, 2007a; 2007b), streamflow in the Lahaina District was used extensively for the cultivation of taro and other agricultural crops by Hawaiian communities. Kumu Pono Associates (2007a) provided some context:

The natural stream alignments were modified and extended in ancient times, with large and small *'auwai* (irrigation channels) constructed to feed thousands of *lo'i kalo* (taro pond fields) in which the primary food crop of the Lahaina region residents was grown...


At least two notable *'auwai*, which span several ahupua'a between Kaua'ula and Kahoma, are described in native lore, and are roughly datable by the chiefs associated with them. The earliest *'auwai* in the Lahainaluna vicinity, is known as *'Auwaiawao*, and is reported named for the Chiefess Wao, sister of Kaululā'au, who ruled a portion of Maui in ca. 1390. The second *'auwai* is known as *'Auwai o Pi'ilani*, and is reported associated with the King, Pi'ilani, who ruled Maui and the neighboring islands in ca. 1450...and has its headwaters in the Kaua'ula Stream.

In 1842, the ancient *'auwai* system extending from Kaua'ula to Lahainaluna was being modified into the Lahainaluna ditch. The ditch was completed in 1847, to facilitate the planting and instructional efforts of Lahainaluna School.

By the 1850s...To develop the sugar plantations...water began to be diverted from *lo'i kalo* to sugar fields...and by the 1860s...diminishing supplies of water were impacting native residency. Indeed, it was reported that in areas where water once flowed, none could then be found.

Individual cultural resources of Kahoma Hydrologic Unit were not classified by the Hawaii Stream Assessment (HSA), but generally classified based on the Historic Preservation Division database. Data were collected in three general areas of: 1) archaeological; 2) historical; and 3) modern practices. Archaeological data were originally compiled by the State Historic Preservation Division and are only current to 1990, the date of the HSA (Table 12-4). The Kahoma hydrologic unit has extensive archeological evidence addressing broad patterns in prehistory, with culturally significant sites including burials, religious structures, and trails. The terracing and *auwai* system that supported *loi kalo* along Kahoma and Kanaha streams are still evident in some locations.

Figure 12-3. Traditional ahupuaa boundaries in the Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Office of Planning, 2015j)



Fishponds

Fishponds are another integral part of traditional Hawaiian culture, which speaks volumes of native Hawaiian skill and knowledge of aquaculture, which has also seen a resurgence of interest in recent years. Fishponds are found throughout the Hawaiian Islands and were either man-made or natural enclosures of water used for the raising and harvesting of fish and other aquatic organisms. Kikuchi (1973) identified six main types of fishponds, two of which are associated with streams (*loko wai*, *loko ia kalo*) and one type is associated with fresh water springs (*kaheka* or *hapunapuna*).

- Type III – *Loko Wai*: An inland fresh water fishpond which is usually either a natural lake or swamp, which can contain ditches connected to a river, stream, or the sea, and which can contain sluice grates. Although most frequently occurring inland, *loko wai* are also located along the coast near the outlet of a stream.
- Type IV – *Loko Ia Kalo*: A fishpond utilizing irrigated taro plots. *Loko ia kalo* are located inland along streams and on the coast in deltas and marshes.
- Type VI – *Kaheka* and *Hapunapuna*: A natural pool or holding pond. The majority, if not all of these types of ponds, are anchialine ponds with naturally occurring shrimp and mollusks.


According to a 1990 Hawaii Coastal Zone Management Program *Hawaiian Fishpond Study for the Islands of Hawaii, Maui, Lanai, and Kauai*, there are no fishponds present in the Kahoma hydrologic unit (DHM, Inc., 1990).

Table 12-3. Cultural resource elements evaluated as part of the Hawaii Stream Assessment for Kahoma hydrologic unit, Maui.

Category	Value
<p>Survey coverage: The extent of archaeological survey coverage was analyzed and recorded as complete, partial, very limited, and none. Few valleys are completely surveyed. Many have little or no survey coverage.</p>	Partial
<p>Predictability: The ability to predict what historic sites might be in unsurveyed areas was scored as high, medium, or low predictability or unable to predict. A high score was assigned if archaeologists were able to predict likely site patterns in a valley given historic documents, extensive archaeological surveys in nearby or similar valleys, and/or partial survey coverage. A low score was assigned if archaeologists were unable to predict site patterns in a valley because of a lack of historical or archaeological information. A medium score was assigned to all other cases.</p>	All others
<p>Number of Sites: The actual number of historic sites known in each valley is straightforward yet very time consuming to count. Instead, archaeologists used survey information to estimate the number of sites in each valley. These figures, adequate for this broad-based assessment, are only rough estimates.</p>	3
<p>Valley significance as a Whole District: The overall evaluation of each valley's significance was made considering each valley a district. The significance criteria of the National and Hawaii Registers of Historic Places were used. Criterion A applies if the district is significant in addressing broad patterns of prehistory or early history. Criterion B applies if the district is associated with important people (rulers) or deities. Criterion C applies if the district contains excellent examples of site types. Criterion D applies if the district is significant for information contained in its sites. Finally, Criterion E applies if the district is culturally significant for traditionally known places or events or for sites such as burials, religious structures, trails, and other culturally noteworthy sites.</p>	ADE
<p>Site Density: The density patterns of historic sites make up a variable extremely important to planners. Three ranks were assigned: low for very few sites due either to normal site patterning or extensive land alteration, moderate for scattered clusters of sites, and high for continuous sites. Valleys with moderate or high density patterns are generally considered moderate or high sensitivity areas.</p>	Moderate
<p>Overall Sensitivity: The overall sensitivity of a valley was ranked very high, high, moderate, low, or unknown. Very high sensitivity areas have moderate or high densities of sites with little or no land alteration. They are extremely important archaeological and/or cultural areas. High sensitivity areas have moderate or high densities of sites with little or no land alteration. Moderate sensitivity areas have very few sites with the sites meriting preservation consideration due to multiple criteria or moderate densities of sites with moderate land alteration. Low sensitivity areas have very few sites due to normal site patterning or due to extensive land alteration. The sites present are significant solely for their informational content, which enable mitigation through data recovery. Those valleys where no surveying had been undertaken and the ability to predict what might be found was low were ranked unknown.</p>	Moderate
<p>Historic Resources: Several types of sites were considered by inclusion in this section, particularly bridges, sugar mills and irrigation systems. Those that are listed on the State or National register were inventoried, but none of them assessed.</p>	No
<p>Taro Cultivation: Streams and stream water have been and continue to be an integral part of the Hawaiian lifestyle. The committee identified a number of factors important to current Hawaiian practices. These include current taro cultivation, the potential for taro cultivation, appurtenant rights, subsistence gathering areas, and stream-related mythology. The committee felt that a complete assessment of the cultural resources of Hawaii's streams should include these items but, due to limits of information, only the current cultivation of taro was included.</p>	No ¹

¹Taro was re-established in Kahoma Valley in 2017 by a community organization, with the first harvest in 2018

Figure 12-4. Tax map key boundaries in the Kahoma hydrologic unit with specific parcels along Kanaha Stream (insert), Maui.



13.0 Public Trust Uses of Water

The State Water Code (Hawaii Revised Statutes 174C-2) states that:

The state water code shall be liberally interpreted to obtain maximum beneficial use of the waters of the State for purposes such as domestic uses, aquaculture uses, irrigation and other agricultural uses, power development, and commercial and industrial uses. However, adequate provision shall be made for the protection of traditional and customary Hawaiian rights, the protection and procreation of fish and wildlife, the maintenance of proper ecological balance and scenic beauty, and the preservation and enhancement of waters of the State for municipal uses, public recreation, public water supply, agriculture, and navigation.

Article 11, Section 1 of the Hawaii State Constitution maintains that the:

State and its political subdivisions shall conserve and protect Hawaii's natural beauty and all natural resources, including land, water, air, minerals, and energy sources, and shall promote the development and utilization of these resources in a manner consistent with their conservation and in furtherance of the self-sufficiency of the State. All public natural resources are held in trust by the State for the benefit of the people.

This solidified the Public Trust Doctrine as constitutional law. Further, Article 11, Section 7, states that the "State has an obligation to protect, control, and regulate the use of Hawaii's water resources for the benefit of its people." The Public Trust Doctrine identifies four priority uses of water as: (1) water for traditional and customary practices, including the growing of taro; (2) reservations of water for Hawaiian Home Land allotments; (3) water for domestic use of the general public; (4) maintenance of waters in its natural state.

In the Kahoma hydrologic unit, the use of water for traditional and customary practices was covered in Chapter 12 and water in its natural state is covered in Chapters 3-7. At a Maui County Council Water Resources Committee in 2017, native Hawaiian cultural practitioners requested that the county refrain from issuing additional water permits until water was restored in Kanaha to support traditional and customary practices. The following is an analysis of reservations of water for Hawaiian Home Lands and domestic water use in Kahoma.


Hawaiian Home Lands

A component in the assessment of water use includes an analysis of the presence of Department of Hawaiian Home Lands (DHHL) parcels within the surface water hydrologic unit. The mission of DHHL is to effectively manage the Hawaiian Home Lands trust and to develop and deliver land to native Hawaiians (PBR Hawaii, 2004). In June 2004, DHHL published the Maui Island Plan which served to examine infrastructure needs, provide development cost estimates, and identify priority areas for homestead development. Of the more than 31,000 acres of DHHL land on the island of Maui, no parcels occurred within the Kahoma hydrologic unit in 2004. However, land in Kahoma originally slated for development by the State of Hawaii Housing and Finance Development Corporation (Leialii Village I-A and I-B) was transferred to DHHL in 2004. The first development (Village I-A) was completed in and beneficiaries began occupying homes in April 2007. The second development (Village I-B) was expected to begin construction in 2015. DHHL now manages two large tracks of land in West Maui that may be served by nearby water resources. The Leialii tract (1-A and 1-B) encompass approximately 75 acres designated for residential development north of Lahaina. By 2031 the estimated potable water demand by DHHL for the Lahaina Aquifer Sector is 0.770 mgd and the non-potable water demand for Honokowai is

2.081 mgd (State of Hawaii, Department of Hawaiian Homelands, 2017). DHHL developed a potable water well mauka of the Honokowai tract that will support the Maui Department of Water Supply Lahaina-Napili Water System. The DHHL groundwater reservation for Honokowai is 0.770 mgd.

The Honokowai track consists of 777 acres of former sugar cane and pineapple fields mauka of the Honoapiilani Highway in Kaanapali, south of the West Maui airport (Figure 13-1). This land was designated for mixed development, with residential, community, supplemental agricultural, general agriculture, commercial and industrial uses. This development was planned before the Leialii tract was designated. As such, there is some uncertainty as to the future development plans with the Leialii tract development getting prioritized over the Honokowai tract. Non-potable water demand for irrigation of supplemental agriculture and general agriculture land use may be met with R-1 recycled water available from the Lahaina Wastewater Treatment Facility (WWTF) or water from Honokohau ditch. The Lahaina WWTF is already connected by a 20-inch line to the Maui Land & Pineapple reservoir at the 300 ft elevation and the Lower Field 14 Reservoir at the 750 ft elevation.

Figure 13-1. Hawaiian Home Lands development parcels and other major land owners, West Maui. (Source: State of Hawaii, Department of Hawaiian Home Lands, 2011)




Domestic Water Supply

Maui County Department of Water Supply operates a surface water treatment facility in Lahaina that utilizes water diverted from Kanaha Stream through the Lahainaluna Ditch. Following treatment, water is mixed with groundwater pumped from Kanaha Well 1 and Kanaha Well 2, however, excess pumping from these wells results in a rise in chloride levels. Surface water is also processed from the Mahinahina Surface Water Treatment Facility from Honokohau Ditch for this system. Napili Wells A, B, and C, and Honokohau Well A are distributed throughout the Maui County Lahaina-Napili system service area and also contribute to the system (Figure 13-2). The Lahaina-Napili water system service area supports much of the single-family connections in the region (Table 13-1).

Table 13-1. Current total and domestic (single family) water use by subdistrict for the Lahaina-Napili service area provided by the Maui County Department of Water Supply. (Source: Maui County, 2018)

subdistrict	Total Use (gpd)	Total Use (mgd)	Single Family Use (gpd)	Single Family % of Total
Lahaina	2,355,746	2.356	435,098	18.5%
Honokowai	2,022,400	2.022	179,658	8.8%
Alaekoa	1,007,878	1.008	417,639	41.4%
Honokohau	2,378	0.002	2,378	100%
Total	5,388,402	5.388	1,034,773	19.2%

Figure 13-2. Water supply service systems for the Lahaina Aquifer System, Maui. (Source: Maui County, 2018)



14.0 Nonstream Uses

Under the State Water Code, nonstream uses are defined as “water that is diverted or removed from its stream channel...and includes the use of stream water outside of the channel for domestic, agricultural, and industrial purposes.” Article XI, Section 3 of the State Constitution states: “The State shall conserve and protect agricultural lands, promote diversified agriculture, increase agricultural self-sufficiency and assure the availability of agriculturally sustainable lands.” Water is crucial to agriculture and agricultural sustainability. Article XI, Section 3 also states, “Lands identified by the State as important agricultural lands needed to fulfill the purposes above shall not be reclassified by the State or rezoned by its political subdivisions without meeting the standards and criteria established by the legislature and approved by a two-thirds vote of the body responsible for the reclassification or rezoning action. [Add Const Con 1978 and election Nov 7, 1978].” It is the availability of water that allows for the designation of Important Agricultural Lands. The Hawaii Farm Bureau Federation, Hawaii’s largest advocacy organization for general agriculture, states that agriculture is a public trust entity worthy of protection, as demonstrated in its inclusion in the State Constitution. They, on behalf of farmers and ranchers, point to the importance of large-scale agriculture to sustainability and self-sufficiency of our islands, particularly in times of catastrophe when imports are cut off.

In most cases, water is diverted from the stream channel via a physical diversion structure. Diversions take many forms, from small PVC pipes in the stream that remove relatively small amounts of water, to earthen auwai (ditches), hand-built rock walls, and concrete dams that remove relatively larger amounts of water.

Water Leaving the Kahoma Hydrologic Unit in Ditch Systems

Water is most often used away from the stream channel and is not returned; however, as in the case of taro fields, water may be returned to the stream at some point downstream of its use. While the return of surface water to the stream would generally be considered a positive value, this introduces the need to consider water quality variables such as increased temperature, nutrients, and dissolved oxygen, which may impact other instream uses. The Kahoma ditches and Kanaha pipeline were designed to remove water from the Kahoma hydrologic unit without returning it to any of the streams in order to supply irrigation water for sugarcane lands, municipal water, and Lahainaluna High School (Table 14-1). Cu

In addition to the amount of water currently (or potentially) being diverted offstream, the Commission must also consider the diversion structure and the type of use, all of which impact instream uses in different ways. The wide range of diversion structures, as noted above, is what makes regulation of surface water particularly difficult, since one standard method cannot be depended upon for monitoring and measuring flow. The ease of diverting streamflow, whether it be by gravity-flow PVC pipe, pump, or a dug channel, also plays a role in the convenience of diverting surface water and the abundance of illegal, non-permitted diversions.

Upon the enactment of the State Water Code and subsequent adoption of the Hawaii Administrative Rules, the Commission required the registration of all existing stream diversions statewide. The Commission categorized the diversions and filed registrations according to the registrant’s last name or company name. While it is recognized that the ownership and/or lease of many of the properties with diversions has changed since then, the file reference (FILEREF) remains the name of the original registrant file (Table 14-1) with locations are depicted in Figure 14-2. Monthly mean daily flow registered by Maui DWS is depicted in Figure 14-1. Data collected by Commission staff site visits, and information extracted from the original registration files regarding the registered diversions may be found in Table 14-2. Mean daily ditch flows diverted from Kahoma and Kanaha streams provided by Pioneer Mill from 1989 to 1994 are provided in Figure 14-3.

Since the enactment of HAR Title 13 Chapter 168, stream diversion works permits are required for the construction of new diversions or alteration of existing diversions, with the exception of routine maintenance. These permitted (as opposed to “registered”) diversion works are not part of the Commission’s verification effort, nor have any diversions been permitted in the Kahoma hydrologic unit.

Table 14-1. Surface water registrations for the Kahoma Hydrologic Unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2015e)

FILEREF	Diversion ID	Stream Name	Active?	Notes
PIONEER MILL	951	Kahoma	Yes	Diversion located on land owned by Kamehameha Schools Bishop Estate. Most streamflow restored in 2010 following discussions with community members. Very little water diverted and transmitted to Kahoma Ranch, which operates some agriculture along with ecotours (ATV, zipline, swimming)
PIONEER MILL	958	Kahoma	No	Diversion abandoned by Pioneer Mill
PIONEER MILL	954	Kanaha	Yes	Diversion originally operated by Pioneer Mill in cooperation with Maui County and Lahainaluna School. Maui County now operates intake for drinking water supply. Intake co-registered by H. Michel
PIONEER MILL	952	Kanaha	No	Diversion abandoned by Pioneer Mill
MICHEL, HANS	853	Kanaha	Yes	Diversion structure is same as 954
MICHEL, HANS	854	Kanaha	Yes	Diversion structure is same as 954
MAUI DWS		Kanaha	Yes	Diversion structure is same as 954; operator of Diversion 954
LAHAINALUNA HS		Kanaha	No	Registered control box that distributed water to Pioneer Mill, Maui DWS and LHS from pipeline drawing Kanaha Stream water; LHS original use was for domestic and agriculture—potable needs met by Maui DWS and agricultural program is currently minimal

Figure 14-1. Registered monthly average mean daily flow (million gallons per day, mgd) diverted from Kanaha Stream in the Kahoma Hydrologic Unit, Maui. (Source: State of Hawaii, Commission on Water Resource Management, 2018c)




Table 14-2. Registered diversions in the Kahoma hydrologic unit.

[Source of photos are denoted at the end of each description; CWRM, Commission on Water Resource Management; Arrows (\Rightarrow) indicate general direction of natural water flow to and out of diversions; Chevrons (\rightrightarrows) indicate direction of diverted surface water flow]

Event ID	File Reference	Tax Map Key	Diversion Amount (cfs)	Active (Yes/No)	Verified (Yes/No)	Riparian (Yes/No)	Rights Claim (Yes/No)
REG.951.6	Pioneer Mill	245022001	0.50	Yes	Yes	No	No

Kahoma Stream is diverted by a concrete dam across the channel that captures approximately 75% of base flow. Water is diverted along the right back into the Kahoma Ditch. There is significant leakage of water over the diversion and past intake pipe through the sluice gate.

Photos. a) Water flowing into the diversion structure from Kahoma Stream and over the diversion (CWRM, 07/2016); b) Diversion structure across the channel (CWRM, 07/2016); c) Diverted water along right bank of stream with intake pipe (CWRM, 07/2016); d) View of ditch and intake pipe with sluice gate (CWRM, 07/2016).



Table 14-2. [Continued]. Registered diversions in the Kahoma hydrologic unit, Maui

Event ID	File Reference	Tax Map Key	Diversion Amount (cfs)	Active (Yes/No)	Verified (Yes/No)	Riparian (Yes/No)	Rights Claim (Yes/No)
REG.954.6	Pioneer Mill	246018007	3.0	Yes	Yes	No	No

Kanaha Stream is diverted by a concrete dam across the channel that captures 100% of base flow. Water is diverted along the left bank into the Lahainaluna Ditch which is piped to a distribution box. There is a small amount of leakage underneath the diversion and along the left bank.

Photos. a) Water flowing into diversion structure from Kanaha Stream (CWRM, 03/2016); b) Diverted water flowing through debris screen in ditch (CWRM, 03/2016); c) Downstream view of leakage from underneath dam (CWRM, 03/2016); d) Upstream view of Kanaha Stream from diversion structure (CWRM, 03/2016).

a)



b)



c)



d)



Table 14-2. [Continued]. Registered diversions in the Kahoma hydrologic unit, Maui

Event ID	File Reference	Tax Map Key	Diversion Amount (cfs)	Active (Yes/No)	Verified (Yes/No)	Riparian (Yes/No)	Rights Claim (Yes/No)
REG.952.6	Pioneer Mill	246018012	0	No	Yes	No	No

Photos. a) View of inactive diversion 952 on Kanaha Stream above confluence with Kahoma Stream (CWRM 03/2018);
a) b)



Table 14-2. [Continued]. Registered diversions in the Kahoma hydrologic unit, Maui

Event ID	File Reference	Tax Map Key	Diversion Amount (cfs)	Active (Yes/No)	Verified (Yes/No)	Riparian (Yes/No)	Rights Claim (Yes/No)
	n/a	246018012	0	No	Yes	No	No


Photos. a) View of new loi intake on Kahoma Stream (CWRM 08/2018); upstream view of Kahoma Stream from intake
a) b)



Table 14-3. Selected off-stream water use statistics in the Kahoma hydrologic unit, Maui, Hawaii. (Source: CWRM, 2018b)

system	stream	operator	Period of record	mean daily flow (mgd)	median flow (mgd)
Kahoma Tunnel	Kahoma	Pioneer Mill	1989-1994	5.59	3.75
Lahainaluna Ditch	Kanaha	Pioneer Mill	1989-1994	2.49	2.20
Lahaina Water Treatment Facility	Kanaha	Maui County	2015-2018	1.76	1.75

Figure 14-3. Mean daily flow in Kahoma Ditch from Kahoma Stream (A) and Lahainaluna Ditch from Kanaha Stream (B) in the Kahoma hydrologic unit. (Source: State of Hawaii, Commission on Water Resource Management, 2018c)



Modifications of Ditch Systems and Groundwater Recharge


Following the establishment of instream flow standards, one of the proposed measures to increase streamflow may be to decrease the amount of water diverted from streams. Such a measure has important implications to ground water recharge because it affects the amount of water available for irrigation. The effects of irrigation water on ground water recharge can be analyzed using the water budget equation¹. Engott and Vana (2007) at the USGS conducted a study that estimated each of the water budget components for west and central Maui using data from 1926 to 2004. Components of the water budget include rainfall, fog drip, irrigation, runoff, evapotranspiration, and recharge. Results of the study were separated into six historical periods: 1926-79, 1980-84, 1985-89, 1990-94, 1995-99, and 2000-04. From 1979 to 2004, ground water recharge decreased 44 percent from 693 million gallons per day to 391 million gallons per day (Figure 14-4). The low recharge rate in 2004 coincides with the lowest irrigation and rainfall rates that were 46 percent and 11 percent lower than those in 1979, respectively. During this

¹ Water-budget is a balance between the amount of water leaving, entering, and being stored in the plant-soil system. The water budget method/equation is often used to estimate ground water recharge.

period, agricultural lands decreased 21 percent from 112,657 acres in 1979 to 88,847 acres in 2004. Further analysis revealed that a 20 percent decrease in irrigation rate could result in a 9 percent reduction in recharge. A similar study by Izuka et al. (2005) reported that a 34 percent decrease in irrigation rate constituted a 7 percent reduction in recharge in the Lihue basin in Kauai, Hawaii.

Droughts, or periods of lower than average rainfall, have been shown to drastically decrease ground water recharge. The period of drought that occurred in 1998-2002, during which rainfall was at least 30 percent lower than the average annual rainfall was estimated to reduce recharge by 27 percent in west and central Maui (Engott and Vana, 2007). In the island of Kauai, the drought conditions reduced recharge in Lihue basin by 34-37 percent (Izuka et al., 2005). Even though droughts can have exacerbating effects on ground water recharge, these effects are transient and are usually mitigated by periods of higher than average rainfall (Engott and Vana, 2007). However, prolonged loss in irrigation water caused by decrease in the amount of water diverted at the ditches has greater effects on the long term trends of ground water levels.

Figure 14-4. Estimated recharge for six historical periods between 1926 and 2004, central and west Maui, Hawaii. (Source: Engott and Vana, 2007)



Utilization of Important Agricultural Lands

The Agricultural Lands of Importance to the State of Hawaii (ALISH) were completed by the State Department of Agriculture (HDOA) in 1977, with the assistance of the Soil Conservation Service (SCS), U.S. Department of Agriculture, and the College of Tropical Agriculture, University of Hawaii. Three classes of agriculturally important lands were established for Hawaii in conjunction with the SCS in an effort to inventory prime agricultural lands nationwide (Figure 14-5). Hawaii's effort resulted in the classification system of lands as: 1) Prime agricultural land; 2) Unique agricultural land; and 3) Other important agricultural land. Each classification was based on specific criteria such as soil characteristics, slope, flood frequency, and water supply. ALISH was intended to serve as a long-term planning guidance for land use decisions related to important agricultural lands. HDOA is currently in the process of developing agricultural incentives based on classifications of Important Agricultural Lands. Kahoma is

comprised of nearly 53 percent of designated agricultural land. Decreasing the amount of water diverted at the ditches located affects the amount of water available for the irrigation of crops on these important agricultural lands.

Table 14-4. Agricultural Lands of Importance to the State of Hawaii and area distributions in the Kahoma hydrologic unit. (Source: State of Hawaii, Office of Planning, 2015g)

Density	Area (mi ²)	Percent of Unit
Unclassified agricultural land	0.107	1.9
Prime agricultural land	1.766	30.8
Other lands	1.138	19.8

As agricultural commodities changed substantially with the closure of largescale pineapple and sugarcane in the 1980s-2000s, the HDOA funded an updated baseline study of agricultural land use for 2015 (Figure 14-6).

Irrigation Needs of the Kahoma Hydrologic Unit

The Commission funded the development of a GIS-based software program the utilizes the state of Irrigation Water Requirement Estimation Decision Support System, IWREDSS (State of Hawaii, Commission on Water Resource Management, 2015b) was developed by the College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa for the State of Hawaii. IWREDSS is an ArcGIS-based numerical simulation model that estimates irrigation requirements (IRR) and water budget components for different crops grown in the Hawaiian environment. The model accounts for different irrigation application systems (e.g., drip, sprinkler, flood), and water application practices (e.g., field capacity versus fixed depth). Model input parameters include rainfall, evaporation, soil water holding capacities, depth of water table, and various crop water management parameters including length of growing season, crop coefficient², rooting depth, and crop evapotranspiration.

Calibration and validation of the model was based on the crop water requirement data for different crops from the Hawaii region United States Department of Agriculture (USDA), Natural Resource Conservation Service (NRCS) Handbook 38 (NRCS-USDA, 1996). Relative errors between the net irrigation requirements (NIR) estimated by the model and those estimated by NRCS range from less than 1 percent to a 26 percent overestimate. This difference may be attributed to the general nature of the technique NRCS used in estimating NIR. Results of the regression analysis indicate a good correlation ($R^2 = 0.97$) between the two techniques; however, the NIR calculations by NRCS were consistently 8 percent higher than those of the IWREDSS model. Overall, the model is an appropriate and practical tool that can be used to assess the IRR of crops in Hawaii.

Understanding that water demand is highly site, weather, application, and crop dependent, IWREDSS can still provide a useful approximation of water needs. According to the simulation results, the IRR for citrus trees is 4000-4500 gallons per acre per day (Table 14-5). The model calculates IRR based on long-term rainfall records available at the weather stations located nearest to the fields. Thus, the estimated IRR represents an average value for given drought scenarios as opposed to average or wet year conditions. However, the estimated IRR for the relative drought year frequencies could be extrapolated to represent the highest demand scenarios. Alternatively, water demand per tree can be used based on the number of trees planted.

² Crop coefficient is an empirically derived dimensionless number that relates potential evapotranspiration to the crop evapotranspiration. The coefficient is crop-specific.

The Kahoma Ranch Reforestation Plan was developed to grow mixed forest, hardwood species, and fruit orchards on former sugar cane lands. The plan calls for planting 120 mango trees, 120 mixed orchard (citrus) trees, 3 acres of mixed dryland and orchard species, and 3 acres of mixed forest and hardwood species. Water is also used for dust control on 3 acres of road surface. At total implementation (2020+), an estimated 113,000 gallons per day will be needed to meet offstream these demands. In 2018, only about 65,000 gallons per day are needed.

Table 14-5. Mean drip irrigation water demand estimates for three crops in the Kahoma service area based on IWREDSS scenarios modeled for two row crop management and two soil management techniques given a 50 ft depth to water table. Demand is based on various drought scenarios.

Crops	row crop management	soil management	estimated irrigation demand (gallons/acre/day) for drought frequency			
			1 in 2 (20%)	1 in 5 (20%)	1 in 10 (10%)	1 in 20 (5%)
citrus	straight row	Poor	4105	4351	4459	4541
	straight row	Good	4090	4338	4448	4530
	contoured	Poor	4088	4336	4446	4528
	contoured	Good	4083	4330	4439	4521
avocado	straight row	Poor	4814	5218	5400	5538
	straight row	Good	4760	5172	5357	5498
	contoured	Poor	4755	5167	5353	5494
	contoured	Good	4718	5137	5327	5471
coffee	straight row	Poor	4640	5006	5171	5295
	straight row	Good	4588	4963	5132	5260
	contoured	Poor	4580	4958	5128	5257
	contoured	Good	4550	4932	5104	5235

Figure 14-5. Agricultural Lands of Importance designated in the Kahoma hydrologic unit, Maui. (Source: State of Hawaii, Office of Planning, 2015g)





Figure 14-6. 2015 Baseline Agricultural Land Use Map for the Kahoma hydrologic Unit, Maui. (Source: (Perroy et al. 2016)



15.0 Bibliography

- Amear, T., Chisholm, I., Beecher, H., Locke, A., and 12 other authors. (2004). Instream flows for riverine resource stewardship, revised edition. Instream Flow Council, Cheyenne, WY, 268 p.
- Anthony, S.S., Hunt, Jr., C.D., Brasher, A.M.D., Miller, L.D., and Tomlinson, M.S. (2004). Water quality on the Island of Oahu, Hawaii, 1999-2001, U.S. Geological Survey Circular 1239, 37 p.
- Coral Reef Assessment and Monitoring Program. (2007). Maui Watershed Information. Hawaii Institute of Marine Biology, Hawaii Coral Reef Assessment and Monitoring Program. Retrieved January 2008, from http://cramp.wcc.hawaii.edu/Watershed_Files/maui/WS_Maui.htm.
- Cowardin, L.M., Carter, V., Golet, F.C., and LaRoe, E.T. (1979). Classification of wetlands and deeper habitats of the United States, U.S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-79/31, 131 p.
- DHM, Inc., Bishop Museum, Public Archaeology Section, Applied Research Group, and Moon, O'Connor, Tam and Yuen. (1990). Hawaiian fishpond study, Islands of Hawaii, Maui, Lanai and Kauai: Prepared for State of Hawaii, Office of State Planning, Coastal Zone Management Program, 196 p.
- Ego, K. (1956). Life history of fresh water gobies. Project no. F-4-R. Fresh Water Game Fish Management Research, Department of Land and Natural Resources, Territory of Hawai'i, Honolulu, 24 p.
- Ekern, P.C., and Chang, J-H. (1985). Pan evaporation: State of Hawaii, 1894-1983. Hawaii Department of Land and Natural Resources, Division of Water and Land Development, Report R74, p.1-3, 38-48.
- Engott, J.A., and Vana, T.T. (2007). Effects of Agricultural Land-Use Changes and Rainfall on Ground-Water Recharge in Central and West Maui, Hawai'i, 1926-2004: U.S. Geological Survey Scientific Investigations Report 2007-5103, 56 p.
- Englund, R., and Filbert, R. (1997). Discovery of the native stream goby, *Lentipes concolor*, above Hawaii's highest waterfall, Hiilawe Falls. Bishop Museum Occasional Papers, v. 49, p. 62-64.
- Englund, R., Polhemus, D. A., and Preston, D. J. (2000). Assessment of the impacts of rainbow trout predation on native aquatic invertebrate species within Kokee State Park streams, Kauai, Hawaii. Bishop Museum Technical Report 18.
- Federal Emergency Management Agency. (2014). FEMA Flood Hazard Zones. Retrieved August 2015, from Hawaii State GIS Web site: http://hawaii.gov/dbedt/gis/data/dfirm_metadata.htm
- Fitzsimons, J.M., and Nishimoto, R.T. (1990). Territories and site tenacity in males of the Hawaiian stream goby *Lentipes concolor* (Pisces: Gobiidae). Ichthyological Exploration of Freshwaters, v. 1, p. 185-189.
- Giambelluca, T.W., Nullet, M.A., Ridgley, M.A., Eyre, P.R., Moncur, J.E.T., and Price, P. (1991). Drought in Hawaii. State of Hawaii, Department of Land and Natural Resources, Division of Water and Land Development, Report R88, 232 p.

- Giambelluca, T.W., and Nullet, D. (1992). Influence of the trade-wind inversion on the climate of a leeward mountain slope in Hawaii. *Climate Research*, 1, p. 207-2126.
- Giambelluca, T.W., Chen, Q., Frazier, A.G., Price, J.P., Chen, Y.-L., Chu, P.-S. Eischeid, J.K., and Delparte D.M. (2013). Online Rainfall Atlas of Hawaii. *Bulletin of the American Meteorological Society*, 94, 313-316.
- Giambelluca, T.W., Shuai, X., Barnes, M.L., Alliss, R.J., Longman, R.J., Miura, T., Chen, Q., Frazier, A.G., Mudd, R.G., Cuo, L., Businger, A.D. (2014). Evapotranspiration of Hawai'i. Final report submitted to the U.S. Army Corps of Engineers—Honolulu District, and the Commission on Water Resource Management, State of Hawai'i.
- Gingerich, S.B. (1999a). Ground water and surface water in the Haiku area, East Maui, Hawaii: U.S. Geological Survey Water-Resources Investigations Report 98-4142, 29 p.
- Gingerich, S.B. (1999b). Ground-water occurrence and contribution to streamflow, Northeast Maui, Hawaii: U.S. Geological Survey Water-Resources Investigations Report 99-4090, 70 p.
- Gingerich, S.B. and Oki, D.S. (2000). Ground Water in Hawaii. U.S. Geological Survey Fact Sheet 126-00.
- Gingerich, S.B. (2005). Median and Low-Flow Characteristics for Streams under Natural and Diverted Conditions, Northeast Maui, Hawaii. U.S. Geological Survey, Scientific Investigations Report 2004-5262, 72 p.
- Gingerich, S.B., Yeung, C.W., Ibarra, T.N., and Engott, J.A. (2007). Water use in wetland kalo cultivation in Hawai'i: U.S. Geological Survey Open-File Report 2007-1157, 68 p. [<http://pubs.usgs.gov/of/2007/1157/>]. Version 1.0, July 24, 2007, Revised Figure 36, p.
- Gingerich, S.B., Engott, J.A. (2012). Groundwater availability in the Lahaina District, West Maui, Hawaii. U.S. Geological Survey, Scientific Investigations Report, 2012-5010, 90 p.
- Grady, C. E. (2006). Volume and Petrologic Characteristics of the Koloa Volcanics, Kauai, Hawaii. Master of Science in Geology and Geophysics. University of Hawaii Manoa
- Handy, E.S.C., Handy, E.G., and Pukui, M.K. (1972). Native Planters in Old Hawaii: Their Life, Lore, and Environment. Bishop Museum Press, Honolulu, Hawaii: Bernice P. Bishop Museum Bulletin 233, 676 p.
- Izuka, S.K., Oki, D.S., and Chen, C. (2005). Effects of irrigation and rainfall reduction on ground-water recharge in the Lihue Basin, Kauai, Hawaii: U.S. Geological Survey Scientific Investigations Report 2005-5146, 48 p.
- Jacobi, J.D. (1989). Vegetation Maps of the Upland Plant Communities on the Islands of Hawai'i, Maui, Moloka'i, and Lana'i, Technical Report Number 68. University of Hawaii at Manoa, Honolulu, Hawaii: Cooperative National Park Resources Studies Unit.
- Juvik, J.O., and Nullet, D. (1995). Relationships between rainfall, cloud-water interception, and canopy throughfall in a Hawaiian montane forest. In Hamilton, L.S., Juvik, J.O., and Scatena, F.N., (eds.), *Tropical montane cloud forests*: New York, Springer-Verlag, p. 165-182.

- Kekaha Agriculture Association (2015). Response to Commission's May 11, 2015 Letter. Schlack Ito LLC, 24 July 2015.
- Kaiser, B., Krause, N., Mecham, D., Wooley, J., and Roumasset, J. (n.d.). Environmental valuation and the Hawaiian economy: Introduction and executive summary, 140 p. Retrieved January 2008, from <http://www.uhero.hawaii.edu/workingpaper/HawaiiEnviroEvaluation.pdf>
- Kido, M. (1996). Recovery processes in Hawaiian streams, p. 76-93. In: Will stream restoration benefit freshwater, estuarine, and marine fisheries?: Proceedings of the October, 1994 Hawaii stream restoration symposium. State of Hawaii Department of Land and Natural Resources, Division of Aquatic Resources, Honolulu.
- Kikuchi, W.K. (1973). Hawaiian aquacultural system. Doctoral thesis, University of Arizona. Manuscript on file in the Bernice P. Bishop Museum Library, Honolulu
- Kumu Pono Associates, LLC. (2007a). A Collection of Traditions and Historical Accounts of Kauaula and Other Lands of Lahaina, Maui, Volume 1 (Part 1). Prepared by Kepa Maly and Onaona Maly. MaKaua111a (060107). pp. Pages ii to 654.
- Kumu Pono Associates, LLC. (2007b). A Collection of Traditions and Historical Accounts of Kauaula and Other Lands of Lahaina, Maui, Volume 1 (Part 2). Prepared by Kepa Maly and Onaona Maly. MaKaua111a (060107). pp. Pages 655 to 1260.
- Lau, L. S. and Mink, J. F. (2006). Hydrology of the Hawaiian Islands. Honolulu: University of Hawaii Press, 274 p.
- Macdonald, G. A., Davis, D. A. and Cox, D. C. (1960). Geology and groundwater resources of the island of Kauai, Hawaii. Haw. Div. Hydrog. Bull. 13, 212 p.
- Maui County. (2018). Draft Maui Island Water Use & Development Plan. Retrieved September 2018, from <https://www.mauicounty.gov/2051/Maui-Island-Water-Use-Development-Plan>
- McRae, M.G. (2007). The potential for source-sink population dynamics in Hawaii's amphidromous species, p. 87-98. In: N.L. Evenhuis and J.M. Fitzsimons (ed.), Biology of Hawaiian streams and estuaries: Proceedings of the symposium on the biology of Hawaiian streams and estuaries. Bishop Museum Bulletin in Cultural and Environmental Studies, v. 3, Honolulu.
- National Park Service, Hawaii Cooperative Park Service Unit. (1990). Hawaii Stream Assessment: A Preliminary Appraisal of Hawaii's Stream Resources: Prepared for State of Hawaii, Commission on Water Resource Management, Report R84, 294 p.
- National Park Service. (2007). Haleakala National Park. Retrieved January 2008, from <http://www.nps.gov/hale/>.
- Nishimoto, R.T., and Kuamoo, D.G.K. (1991). The occurrence and distribution of the native goby (*Lentipes concolor*) in Hawai'i Island streams with notes on the distribution of the native goby species, p. 77-95. In: W. Devick (ed.), New directions in research, management and conservation of Hawaiian freshwater stream ecosystems: Proceedings of the 1990 symposium on stream biology and fisheries management. State of Hawaii Department of Land and Natural Resources, Division of Aquatic Resources, Honolulu.

- Nishimoto, R.T., and Kuamoo, D.G.K. (1997). Recruitment of goby postlarvae into Hakalau stream, Hawai'i Island. *Micronesica*, v. 30, p. 41-49.
- Nullet, D. (1987). Energy sources for evaporation on tropical islands. *Physical Geography*, 8: p. 36-45.
- Nullet, D., and Giambelluca, T.W. (1990). Winter evaporation on a mountain slope, Hawaii. *Journal of Hydrology*: 112, p. 257-265.
- Oki, D. (2003). Surface Water in Hawaii. U.S. Geological Survey, Fact Sheet 045-03.
- Oki, D. (2004). Trends in Streamflow Characteristics at Long-Term Gaging Stations, Hawaii. U.S. Geological Survey, Scientific Investigations Report 2004-5080, 120 p.
- OmniTrak Group Inc. (2007). Hawaii State Parks Survey: Prepared for Hawaii Tourism Authority, 98 p. Retrieved February 2008, from http://www.hawaii-tourism-authority.org/documents_upload_path/reports/HTAPRO-Report-12-01-2007.pdf
- Pacific Disaster Center. (2007). Natural Hazards: Flood. Retrieved March 2008, from <http://www.pdc.org/iweb/flood.jsp?subg=1>.
- PBR Hawaii. (2004). Maui Island Plan: Prepared for State of Hawaii, Department of Hawaiian Home Lands, 340 p.
- Penn, D.C. (1997). Water and energy flows in Hawai'i taro pondfields: Honolulu, Hawaii, University of Hawaii at Manoa, Ph.D. dissertation, 376 p.
- Radtke, R.L., Kinzie III, R.A., and Folsom, S.D. (1988). Age at recruitment of Hawaiian freshwater gobies. *Environmental Biology of Fishes*, v. 23, p. 205-213.
- Scholl, M.A., Gingerich, S.B., and Tribble, G.W. (2002). The influence of microclimates and fog on stable isotope signatures used in interpretation of regional hydrology: East Maui, Hawaii. *Journal of Hydrology*, 264 (2002), p. 170-184.
- Scott, J.M., Mountainspring, S., Ramsey, and Kepler, C.B. (1986). Forest Bird Communities of the Hawaiian Islands: Their Dynamics, Ecology and Conservation. *Studies in Avian Biology*, No. 9, 431 p.
- Shade, P.J. (1999). Water budget of East Maui, Hawaii, U.S. Geological Survey Water Resources Investigations Report 98-4159, p. 36.
- Sherrod, D.R., Sinton, J.M., Watkins, S. E., and Brunt, K.M. (2007). Geological Map of the State of Hawaii: U.S. Geological Survey Open-File Report 2007-1089, 83 p., 8 plates, scales 1:100,000 and 1:250,000, with GIS database. U.S. Geological Survey.
- State of Hawaii, Commission on Water Resource Management. (2005a). Commission on Water Resource Management surface-water hydrologic units: A management tool for instream flow standards. Department of Land and Natural Resources, Commission on Water Resource Management, Report PR-2005-01, 111 p.

- State of Hawaii, Commission on Water Resource Management. (2005b). Hawaii Drought Plan 2005 Update. State of Hawaii, Department of Land and Natural Resources, Division of Water and Land Development, p. 1-1 to 5-4.
- State of Hawaii, Commission on Water Resource Management. (2007). Hawaii Water Plan: Water Resource Protection Plan, Public review draft. Department of Land and Natural Resources, Commission on Water Resource Management, 557 p.
- State of Hawaii, Commission on Water Resource Management. (2015a). Irrigation Ditch GIS Layer [Database file]. Retrieved July 2015.
- State of Hawaii, Commission on Water Resource Management. (2015b). Water Management Software to Estimate Crop Irrigation Requirement for Consumptive Use Permitting In Hawaii. Department of Land and Natural Resources, 61 p.
- State of Hawaii, Commission on Water Resource Management. (2015c). Surface-water hydrologic unit GIS data layer. Department of Land and Natural Resources, Commission on Water Resource Management. Retrieved July 2015.
- State of Hawaii, Commission on Water Resource Management. (2015e). Diversion GIS Layer [Database file]. Retrieved July 2015.
- State of Hawaii, Commission on Water Resource Management. (2017). State Water Projects Plan Update: Department of Hawaiian Home Lands. <http://files.hawaii.gov/dlnr/cwrm/planning/swpp2017.pdf>
- State of Hawaii, Commission on Water Resource Management. (2018a). Mahinahina Deep Monitoring Well data. Retrieved June 2018 from <http://dlnr.hawaii.gov/cwrm/groundwater/monitoring/>
- State of Hawaii, Commission on Water Resource Management. (2018b). Well index database [Database file]. Retrieved June 2018.
- State of Hawaii, Commission on Water Resource Management. (2018c). Water Use Reporting [Database file]. Retrieved July 2018.
- State of Hawaii, Department of Hawaiian Homelands. (2011). West Maui (Leialii-Honokowai) Regional Plan.
- State of Hawaii, Department of Health, Environmental Planning Office. (1987). Water Quality Standards Map of the Island of Maui [map]. Retrieved July 2015, from <http://hawaii.gov/health/environmental/water/cleanwater/wqsmaps/index.html>.
- State of Hawaii, Department of Health. (2004). Amendment and Compilation of Chapter 11-54, Hawaii Administrative Rules. Retrieved July 2015 from <http://hawaii.gov/health/about/rules/11-54.pdf>.
- State of Hawaii, Department of Health, Environmental Planning Office. (2001). Hawaii's Water Quality Standards: A Public Guide [brochure]. Retrieved July 2015 from <http://hawaii.gov/health/environmental/env-planning/wqm/wqsbrochure.pdf>.

- State of Hawaii, Department of Health, Environmental Planning Office. (2004). Final 2004 List of Impaired Waters in Hawaii Prepared Under Clean Water Act §303(d). Retrieved January 2008 from <http://hawaii.gov/health/environmental/env-planning/wqm/wqm.html>.
- State of Hawaii, Department of Health, Environmental Planning Office. (2007). 2006 State of Hawaii Water Quality Monitoring and Assessment Report: Integrated Report to the U.S. Environmental Protection Agency and The U.S. Congress Pursuant To Sections §303(D) and §305(B) Clean Water Act (P.L. 97-117). Retrieved January 2008 from <http://hawaii.gov/health/environmental/env-planning/wqm/wqm.html>.
- State of Hawaii, Division of Aquatic Resources. (1993). Native stream animals [Brochure].
- State of Hawaii, Division of Aquatic Resources. (2015). Point-quadrat survey database. Accessed October 2015.
- State of Hawaii, Division of Forestry and Wildlife. (2008a). Hawaii Forest Reserve System. Retrieved July 2015, from <http://www.state.hi.us/dlnr/dofaw/frs/page2.htm>.
- State of Hawaii, Division of Forestry and Wildlife. (2008b). Watershed Partnership Program. Retrieved July 2015, from <http://www.state.hi.us/dlnr/dofaw/wpp/index.html>
- State of Hawaii, Office of Planning. (1996). Stream gauges [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/strmgage.htm>
- State of Hawaii, Office of Planning. (1999). Parks [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/parks.htm>
- State of Hawaii, Office of Planning. (2002a). Elementary school districts [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Data Server.
- State of Hawaii, Office of Planning. (2002b). Hunting areas [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/huntareas.htm>
- State of Hawaii, Office of Planning. (2002c). Na Ala Hele state trails and access system [GIS data file]. Retrieved December 2007, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/nahtrails.htm>
- State of Hawaii, Office of Planning. (2002d). Public school locations [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/pubschools.htm>
- State of Hawaii, Office of Planning. (2003). National wetlands inventory [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/wetlnds.htm>
- State of Hawaii, Office of Planning. (2004a). Coastal resources [GIS data file]. Retrieved December 2007, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/cstrsrc.htm>
- State of Hawaii, Office of Planning. (2004b). Critical habitat [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/criticalhab.htm>

State of Hawaii, Office of Planning. (2004c). Ditches [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/ditches.htm>

State of Hawaii, Office of Planning. (2004d). Vegetation [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/veg.htm>

State of Hawaii, Office of Planning. (2004e). 500 foot contours [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/cntrs500.htm>

State of Hawaii, Office of Planning. (2005). Division of Aquatic Resources (DAR) stream [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/streams.htm>

State of Hawaii, Office of Planning. (2007b). Reserves [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/reserves.htm>

State of Hawaii, Office of Planning. (2008). Conservation District Subzones [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/cdsubzn.htm>

State of Hawaii, Office of Planning. (2015a). Dams [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/dams.htm>

State of Hawaii, Office of Planning. (2015b). Aquifers [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/dlnraq.htm>

State of Hawaii, Office of Planning. (2015c). Solar radiation [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/solarad.htm>

State of Hawaii, Office of Planning. (2015d). State land use district [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/slud.htm>

State of Hawaii, Office of Planning. (2015e). Water Quality Classifications [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/classwater.htm>

State of Hawaii, Office of Planning. (2015f). GAP Land Cover Analysis [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <ftp://ftp.gap.uidaho.edu/products/Hawaii.zip>

State of Hawaii, Office of Planning. (2015g). ALISH [GIS data file]. Retrieved October 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/alish.htm>

State of Hawaii, Office of Planning. (2015h). ALUM [GIS data file]. Retrieved October 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/alum.htm>

State of Hawaii, Office of Planning. (2015i). Areas of threatened and endangered plants [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/reserves.htm>

State of Hawaii, Office of Planning. (2015j). Historic land divisions (Ahupuaa) for the island of Maui [GIS data file]. Retrieved December 2007, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/histlanddiv.htm>

- State of Hawaii, Office of Planning. (2015k). C-CAP Land Cover Analysis: Impervious Surface and Land Cover Data [GIS data file]. National Oceanic and Atmospheric Administration, Coastal Services Center. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <ftp://ftp.gap.uidaho.edu/products/Hawaii.zip>
- State of Hawaii, Office of Planning. (2015l). census blocks [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://files.hawaii.gov/dbedt/op/gis/data/blocks10.htm>
- State of Hawaii, Office of Planning. (2015m). Soils [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/soils.htm>
- State of Hawaii, Office of Planning. (2015n). Wetlands [GIS data file]. Retrieved July 2015, from Hawaii Statewide GIS Program Web site: <http://hawaii.gov/dbedt/gis/wetlands.htm>
- State of Nevada, Department of Conservation and Natural Resources, Division of Water Resources. (n.d.). Water words dictionary: Technical water, water quality, environmental, and water-related terms. Retrieved February 2008, from <http://water.nv.gov/WaterPlanning/dict-1/ww-dictionary.pdf>
- Stearns, H.T. and MacDonald, G.A. (1942). Geology and Ground-Water Resources of the Island of Maui, Hawaii. [Hawaii Division of Hydrography, Bulletin 7]. [In cooperation with the Geological Survey, United States Department of the Interior.]
- Trust for Public Land. (1998). East Maui resource inventory: Prepared for National Park Service, Rivers, Trails, and Conservation Assistance Program, with the assistance of Bay Pacific Consulting, 101 p.
- University of Hawaii. (2003). Drought Risk and Vulnerability Assessment and GIS Mapping Project. State of Hawaii, Department of Land and Natural Resources, Division of Water and Land Development, 157 p.
- U. S. Department of Agriculture, Natural Resources Conservation Service, Conservation Engineering Division. (1986). Appendix A: Hydrologic Soil Groups. Urban Hydrology for Small Watersheds. (Technical Release 55.) Retrieved December 26, 2007 from <http://www.info.usda.gov/CED/ftp/CED/tr55.pdf>
- U.S. Department of Agriculture, Soils Conservation Service. [In cooperation with The University of Hawaii Agriculture Experiment Station.] (1972). Soil Survey of Islands of Kauai, Oahu, Maui, Molokai, and Lanai, State of Hawaii. Washington, DC: U.S. Government Printing Office.
- U.S. Environmental Protection Agency. (2008). "Impaired Waters and Total Maximum Daily Loads." Retrieved April 23, 2008 from <http://www.epa.gov/owow/tmdl/intro.html>
- U.S. Environmental Protection Agency, Region 9, and State of Hawaii, Department of Health. (2002). Revisions to total maximum daily loads for the Ala Wai Canal, Island of Oahu, Hawaii. Retrieved February 2008, from <http://hawaii.gov/health/environmental/env-planning/wqm/awtmdlfinal.pdf>
- U.S. Geological Survey. (1996). Digital raster graphics (DRG) data. Retrieved July 2015, from <http://data.geocomm.com/drg/index.html>.
- U.S. Geological Survey. (2001). Digital elevation model (DEM) – 10 meter. Retrieved July 2015, from <http://data.geocomm.com/dem/>.

U.S. Geological Survey. (2007). Geologic Map of the State of Hawaii. [With GIS database.] (Open File Report 2007-1089). Reston, VA: D.S. Sherrod, J.M. Sinton, S.E. Watkins, and K.M. Brunt.

U.S. Geological Survey. (2011). National Gap Analysis Project Land Cover Data Portal.
<https://gapanalysis.usgs.gov/gaplandcover/data/download/>

Wilcox, C. (1996). Sugar water: Hawaii's plantation ditches: Honolulu, University of Hawaii Press, p. 191.

Wilson Okamoto & Associates, Inc. (1983). Instream Use Study, Windward Oahu: Prepared for State of Hawaii, Division of Water and Land Development, Report R68, 154 p.

Ziegler, A.C. (2002). Hawaiian natural history, ecology, and evolution. Honolulu: University of Hawaii Press, 477 p.

16.0 Appendices


Appendix A Kahoma, Maui, Hawaii. June 2008. DAR Watershed Code: 61005
State of Hawaii, Department of Land and Natural Resources, Division of Aquatic Resources.

APPENDIX A

**State of Hawai'i, Department of Land and Natural Resources,
Division of Aquatic Resources**

**Atlas of Hawaiian Watersheds & Their Aquatic Resources
Kahoma, Maui**

Kahoma, Maui



WATERSHED FEATURES

Kahoma watershed occurs on the island of Maui. The Hawaiian meaning of the name is “the thin one”. The area of the watershed is 5.8 square mi (14.9 square km), with maximum elevation of 5,764 ft (1,757 m). The watershed’s DAR cluster code is 5, meaning that the watershed is medium size, steep in the upper watershed, and with little embayment. The percent of the watershed in the different land use districts is as follows: 14.4% agricultural, 70.7% conservation, 0% rural, and 14.9% urban.

Land Stewardship: Percentage of the land in the watershed managed or controlled by the corresponding agency or entity. Note that this is not necessarily ownership.

<u>Military</u>	<u>Federal</u>	<u>State</u>	<u>OHA</u>	<u>County</u>	<u>Nature Conservancy</u>	<u>Other Private</u>
0.0	0.0	30.0	0.0	0.0	0.1	69.9

Land Management Status: Percentage of the watershed in the categories of biodiversity protection and management created by the Hawaii GAP program.

Permanent Biodiversity <u>Protection</u>	Managed for Multiple <u>Uses</u>	Protected but <u>Unmanaged</u>	<u>Unprotected</u>
0.1	59.4	0.5	40.0

Land Use: Areas of the various categories of land use. These data are based on NOAA C-CAP remote sensing project.

	<u>Percent</u>	<u>Square mi</u>	<u>Square km</u>
High Intensity Developed	1.4	0.08	0.21
Low Intensity Developed	3.4	0.20	0.51
Cultivated	0.9	0.05	0.13
Grassland	12.5	0.72	1.86
Scrub/Shrub	30.3	1.75	4.53
Evergreen Forest	50.4	2.91	7.53
Palustrine Forested	0.0	0.00	0.00
Palustrine Scrub/Shrub	0.0	0.00	0.00
Palustrine Emergent	0.0	0.00	0.00
Estuarine Forested	0.0	0.00	0.00
Bare Land	0.1	0.01	0.02
Unconsolidated Shoreline	0.2	0.01	0.03
Water	0.7	0.04	0.10
Unclassified	0.0	0.00	0.00

STREAM FEATURES

Kahoma is a perennial stream. Total stream length is 17.4 mi (28 km). The terminal stream order is 5.

Reach Type Percentages: The percentage of the stream's channel length in each of the reach type categories.

<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
0.0	1.4	24.8	45.0	28.8

The following stream(s) occur in the watershed:

Hālonā Kahoma Kanahā

BIOTIC SAMPLING EFFORT

Biotic samples were gathered in the following year(s):

1962 1963

Distribution of Biotic Sampling: The number of survey locations that were sampled in the various reach types.

<u>Survey type</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
HDFG	0	1	1	2	0
Reservoir	0	0	1	0	0

BIOTA INFORMATION

Species List

Native Species

Crustaceans *Atyoida bisulcata*
Fish Gobiid sp.

Native Species

Insects Diptera sp.
Megalagrion sp.
Telmatogeton sp.

Introduced Species

Fish *Tilapia sp.*

Species found in Impoundments

Fish *Tilapia sp.*

Species Distributions: Presence (P) of species in different stream reaches.

<u>Scientific Name</u>	<u>Status</u>	<u>Estuary</u>	<u>Lower</u>	<u>Middle</u>	<u>Upper</u>	<u>Headwaters</u>
<i>Atyoida bisulcata</i>	Endemic		P		P	
<i>Megalagrion sp.</i>	Endemic		P	P	P	
Gobiid sp.	Indigenous		P	P		
<i>Telmatogeton sp.</i>	Indigenous				P	
<i>Tilapia sp.</i>	Introduced			P		
Diptera sp.	Undetermined			P		

HISTORIC RANKINGS

Historic Rankings: These are rankings of streams from historical studies. "Yes" means the stream was considered worthy of protection by that method. Some methods include non-biotic data in their determination. See Atlas Key for details.

Multi-Attribute Prioritization of Streams - Potential Heritage Streams (1998): No

Hawaii Stream Assessment Rank (1990): Moderate

U.S. Fish and Wildlife Service High Quality Stream (1988): No

The Nature Conservancy- Priority Aquatic Sites (1985): No

National Park Service - Nationwide Rivers Inventory (1982): No

Current DAR Decision Rule Status: The following criteria are used by DAR to consider the biotic importance of streams. "Yes" means that watershed has that quality.


Native Insect Diversity <u>> 19 spp.</u>	Native Macrofauna <u>Diversity > 5 spp.</u>	Absence of Priority 1 <u>Introduced spp.</u>
No	No	No
Abundance of Any <u>Native Species</u>	Presence of Candidate <u>Endangered Species</u>	Endangered Newcomb's <u>Snail Habitat</u>
No	No	No

CURRENT WATERSHED AND STREAM RATINGS


The current watershed and stream ratings are based on the data contained in the DAR Aquatic Surveys Database. The ratings provide the score for the individual watershed or stream, the distribution of ratings for that island, and the distribution of ratings statewide. This allows a better understanding of the meaning of a particular ranking and how it compares to other streams. The ratings are standardized to range from 0 to 10 (0 is lowest and 10 is highest rating) for each variable and the totals are also standardized so that the rating is not the average of each component rating. These ratings are subject to change as more data are entered into the DAR Aquatic Surveys Database and can be automatically recalculated as the data improve. In addition to the ratings, we have also provided an estimate of the confidence level of the ratings. This is called rating strength. The higher the rating strength the more likely the data and rankings represent the actual condition of the watershed, stream, and aquatic biota.

WATERSHED RATING: Kahoma, Maui


Land Cover Rating: Rating is based on a scoring system where in general forested lands score positively and developed lands score negatively.



Shallow Waters Rating: Rating is based on a combination of the extent of estuarine and shallow marine areas associated with the watershed and stream.




Stewardship Rating: Rating is based on a scoring system where higher levels of land and biodiversity protection within the watershed score positively.




WATERSHED RATING (Cont): Kahoma, Maui


Size Rating: Rating is based on the watershed area and total stream length. Larger watersheds and streams score more positively.




Wetness Rating: Rating is based on the average annual rainfall within the watershed. Higher rainfall totals score more positively.



Reach Diversity Rating: Rating is based on the types and amounts of different stream reaches available in the watershed. More area in different reach types score more positively.



Total Watershed Rating: Rating is based on combination of Land Cover Rating, Shallow Waters Rating, Stewardship Rating, Size Rating, Wetness Rating, and Reach Diversity Rating.





BIOLOGICAL RATING: Kahoma, Maui

Native Species Rating: Rating is based on the number of native species observed in the watershed.

Stream Rating



1



Introduced Genera Rating: Rating is based on the number of introduced genera observed in the watershed.

Stream Rating



9



All Species' Score Rating: Rating is based on the Hawaii Stream Assessment scoring system where native species score positively and introduced species score negatively.

Stream Rating



5



Total Biological Rating: Rating is the combination of the Native Species Rating, Introduced Genera Rating, and the All Species' Score Rating.


Stream Rating

4




OVERALL RATING: Kahoma, Maui

Overall Rating: Rating is a combination of the Total Watershed Rating and the Total Biological Rating.



RATING STRENGTH: Kahoma, Maui

Rating Strength: Represents an estimate of the overall study effort in the stream and is a combination of the number of studies, number of different reaches surveyed, and the number of different survey types.



REFERENCES

- 1963. Shima, S.I. Limnological Survey for Introduction of Exotic Species of Fish.
- 2008. Hawai'i Division of Aquatic Resources. DAR Point Quadrat Survey Data from the DAR Aquatic Surveys Database.
- 2008. Hawai'i Division of Aquatic Resources. Impoundment Surveys in DAR Aquatic Surveys Database.

Blank Page