



See inside for complete **Precautionary Statements, Directions for Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators, and other handlers must wear long-sleeved shirt, long pants, shoes plus socks.

Workers performing cleaning, maintenance, and repair of equipment must wear: coveralls over long-sleeved shirt, long pants, shoes plus socks, and chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, mammals, fish and aquatic invertebrates. Do not contaminate water when disposing of equipment washwater or rinsate. Do not plant treated seeds by broadcasting to the soil surface. Ensure that all seeds are thoroughly covered with soil. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE: Store in original container away from feed and food. Store in cool, dry place. Do not store in direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER HANDLING AND DISPOSAL: Non-refillable container (greater than 5 gallons). Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures allowed by state and local authorities.

Acceleron®, Monsanto and Vine Design® and NemaStrike™ are trademarks of Monsanto Technology LLC.

©2017

For **MEDICAL And TRANSPORTATION Emergencies ONLY** Call Collect 24 Hours A Day: 1-314-694-4000.
For **PRODUCT USE** Information, Call 1-877-775-8787.

EPA Reg. No. 524-624

EPA Est. 5905-GA-01

20170501

PRODUCED FOR:
Monsanto Company
800 N. Lindbergh Blvd.
St. Louis, Missouri 63167

Lot No. _____



ACCELERON® NemaStrike™ ST

Seed treatment for nematode control on
corn, cotton and soybean.

ACTIVE INGREDIENT:

tioxazafen*, 3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole 45.9%

OTHER INGREDIENTS: 54.1%
100.0%

* This product is a suspension concentrate containing 4.51 pounds of tioxazafen per gallon.

EPA Reg. No. 524-624

KEEP OUT OF REACH OF CHILDREN

CAUTION

HOTLINE NUMBER

In case of an emergency involving this product, or for medical assistance,
call collect, day or night, (314) 694-4000.

For product use information regarding Acceleron® Seed Applied Solutions call:
1-877-775-8787

Produced for: Monsanto Company • 800 N. Lindbergh Blvd.
St. Louis, MO 63167 USA

NET 30 GAL

20
CODE

2D
CODE

116004RT-1

OPEN
HERE!

ACCELERON[®] *NemaStrike[™] ST*

Seed treatment for nematode control on
corn, cotton and soybean.

ACTIVE INGREDIENT:

tioxazafen*, 3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole 45.9%

OTHER INGREDIENTS: 54.1%
100.0%

6.625"

* This product is a suspension concentrate containing 4.51 pounds of tioxazafen per gallon.

EPA Reg. No. 524-624

KEEP OUT OF REACH OF CHILDREN

CAUTION

HOTLINE NUMBER

In case of an emergency involving this product, or for medical assistance,
call collect, day or night, (314) 694-4000.

For product use information regarding Acceleron[®] Seed Applied Solutions call:
1-877-775-8787

Produced for: Monsanto Company • 800 N. Lindbergh Blvd.
St. Louis, MO 63167 USA

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children

CAUTION

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators, and other handlers must wear long-sleeved shirt, long pants, shoes plus socks.

Workers performing cleaning, maintenance, and repair of equipment must wear: coveralls over long-sleeved shirt, long pants, shoes plus socks, and chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Environmental Hazards

This product is toxic to birds, mammals, fish and aquatic invertebrates. Do not contaminate water when disposing of equipment washwater or rinsate.

Do not plant treated seeds by broadcasting to the soil surface. Ensure that all seeds are thoroughly covered with soil. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

DIRECTIONS FOR USE

Read the entire label before using this product.

It is a violation of US Federal law to use this product in a manner inconsistent with its labeling.

For use only in commercial seed treatment facilities using fully-automated closed transfer and application equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications.

Restrictions

- Do not treat seeds on farm or in field.
- To be used in commercial seed treatment facilities using fully-automated closed transfer and application equipment ONLY
- Do not plant treated seeds by broadcasting to the soil surface.

Application Information

USE INFORMATION

Apply Acceleron® NemaStrike™ ST as a water-based slurry either alone or with registered seed treatments using fully-automated and closed standard slurry or mist-type seed treatment application equipment and closed transfer equipment.

Acceleron® NemaStrike™ ST may be used in combination with other registered Acceleron® Seed Applied Solutions. Pre-test for compatibility with other seed treatment products. Users must follow the most restrictive directions for use and precautionary statements of each product that is co-applied.

Uniform application of seed is necessary to ensure seed safety and best disease protection. Dilute with a sufficient amount of water to achieve uniform application to seed which is necessary to ensure seed safety and best disease control. Use high quality, properly cleaned seed.

All seed treated with this product must be adequately dyed with a suitable color to prevent accidental use as food for humans or feed for animals. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. Any dye or colorant added to treated seed must be cleared for use pursuant to 40 CFR, Part 153.155.

MIXING INSTRUCTIONS

Add approximately 1/2 to 2/3 of the required water to the treatment slurry tank followed by the accompanying seed treatment products and Acceleron® NemaStrike™ ST. The order of product addition is dependent on the accompanying seed treatment products and application equipment used. Allow each slurry component to disperse completely prior to the next addition until a uniform suspension is obtained. After all slurry components have been added and dispersed completely add the remaining 1/2 to 1/3 of required water for proper slurry volume and mix until uniform. Maintain adequate agitation until treatment slurry is applied.

APPLICATION RATES TO CONTROL NEMATODES

Crop	Use Rates	Maximum Annual Rates
Corn, field	0.5 – 1.0 mg a.i./ seed or 2.5 – 5.0 fl oz product/ 80,000 seeds	Do not apply more than 0.089 lb a.i./acre/year. This is based on a maximum seeding rate of 40,250 seeds/acre or 33.2 lbs seed/acre
Soybean (<i>Glycine</i> spp.)	0.25 – 0.5 mg a.i./ seed or 2.2 – 4.4 fl oz product/ 140,000 seeds	Do not apply more than 0.28 lb a.i./acre/year. This is based on a maximum seeding rate of 250,000 seeds/acre or 166.7 lbs seed/acre
Cotton	0.5 – 1.0 mg a.i./ seed or 7.8 – 15.6 fl oz product/ 250,000 seeds	Do not apply more than 0.19 lb a.i./acre/year. This is based on a maximum seeding rate of 85,000 seeds/acre or 18.9 lbs seed/acre

SEED LABELING

The user is responsible for ensuring that the seed container meets all of the requirements under the Federal Seed Act. The Federal Seed Act requires that the container of seed treated with Acceleron®, NemaStrike™ ST must be labeled with the following statements:

- This seed has been treated with tioxazafen.
- Do not use treated seed for food, feed or oil production.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated Acceleron® NemaStrike™ ST:

- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils when handling treated seed.
- Store treated seed away from food and feed, and, do not allow access by children, pets, or livestock. Do not reuse empty seed bags.
- Surplus treated seed or empty seed containers should be stored or disposed according to local requirements.

Do not contaminate bodies of water when disposing of excess treated seed or wash waters of planting equipment.

Treated seed is toxic to birds and mammals. Exposed seed is hazardous to birds and mammals that come in contact with treated seed. Cover or collect treated seeds that are spilled during loading and planting.

Do not plant seeds by broadcasting to the soil surface.

- For corn and soybeans: Treated seeds must be planted at a minimum depth of 1 inch.
- For cotton: Treated seeds must be planted at a minimum depth of 0.5 inches.

Do not use at a rate that will result in more than 0.28 lb tioxazafen per acre per year.

In case of early season crop failure, seeds (treated or untreated) may be immediately replanted as long as the maximum yearly application rate per acre is not exceeded.

Plant-back Restrictions: Corn, cotton, and soybeans may be replanted immediately. All other crops may be planted after 30 days.

CROP ROTATION

Crops listed on this label may be replanted at any time as long as yearly maximum application rate per acre is not exceeded. All other crops may be planted after 30 days.

In case of early season crop failure, seeds (treated or untreated) may be replanted immediately as long as yearly maximum application rate per acre is not exceeded.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

PESTICIDE STORAGE: Store in original container away from feed and food. Store in a cool, dry place. Do not store in direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER HANDLING AND DISPOSAL: Non-refillable container (5 gallons or less). Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures allowed by state and local authorities.

Non-refillable container (greater than 5 gallons). Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other side and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or by other procedures allowed by state and local authorities.

LIMITS OF LIABILITY AND WARRANTY

Monsanto Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use (“Directions”) when used in accordance with those Directions under the conditions described therein. TO THE EXTENT, CONSISTENT WITH APPLICABLE LAW, NO OTHER, EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, crop injury, ineffectiveness or other unintended consequences that may result because of such factors as weather conditions at planting, environmental conditions during seed storage, presence of other materials, seed of low quality or low vigor or low germination, or the manner of use or application inconsistent with that set for in the Directions.

This Company does not warrant any product, reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY, OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT, CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

20170501

©2017

Produced for:

Monsanto Company

800 N. Lindbergh Blvd.

St. Louis, MO 63167 USA

EPA Reg. No. 524-624

Acceleron®, Monsanto and Vine Design® and
NemaStrike™ are trademarks of Monsanto
Technology LLC.



